

Form PTO-1390 U.S. DEPARTMENT OF COMMERCE PATENT AND TRADEMARK OFFICE (REV 10-95)		ATTORNEY'S DOCKET NUMBER <b>702-001463</b>
TRANSMITTAL LETTER TO THE UNITED STATES DESIGNATED/ELECTED OFFICE (DO/EO/US) CONCERNING A FILING UNDER 35 U.S.C. 371		U.S. APPLICATION NO. (If known, see 37 CFR 1.5) <b>09/601490</b>
INTERNATIONAL APPLICATION NO <b>PCT/EP99/00748</b>	INTERNATIONAL FILING DATE <b>04.02.99 (February 04, 1999)</b>	PRIORITY DATES CLAIMED <b>04.02.98 and 06.02.98</b>
TITLE OF INVENTION <b>IDENTIFICATION, PRODUCTION AND USE OF STAPHYLOKINASE DERIVATIVES WITH REDUCED IMMUNOGENICITY AND/OR REDUCED CLEARANCE</b>		
APPLICANT(S) FOR DO/EO/US <b>Désiré José COLLEN</b>		
<p>Applicant herewith submits to the United States Designated/Elected Office (DO/EO/US) the following items and other information</p> <p>1 <input checked="" type="checkbox"/> This is a <b>FIRST</b> submission of items concerning a filing under 35 U.S.C. 371.</p> <p>2 <input type="checkbox"/> This is a <b>SECOND</b> or <b>SUBSEQUENT</b> submission of items concerning a filing under 35 U.S.C. 371.</p> <p>3 <input checked="" type="checkbox"/> This express request to begin national examination procedures (35 U.S.C. 371(f)) at any time rather than delay examination until the expiration of the applicable time limit set in 35 U.S.C. 371(b) and PCT Articles 22 and 39(1).</p> <p>4 <input checked="" type="checkbox"/> A proper Demand for International Preliminary Examination was made by the 19th month from the earliest claimed priority date.</p> <p>5 <input checked="" type="checkbox"/> A copy of the International Application as filed (35 U.S.C. 371(c)(2))</p> <p>a <input type="checkbox"/> is transmitted herewith (required only if not transmitted by the International Bureau)</p> <p>b <input checked="" type="checkbox"/> has been transmitted by the International Bureau</p> <p>c <input type="checkbox"/> is not required, as the application was filed in the United States Receiving Office (RO/US).</p> <p>6 <input type="checkbox"/> A translation of the International Application into English (35 U.S.C. 371(c)(2)).</p> <p>7 <input checked="" type="checkbox"/> Amendments to the claims of the International Application under PCT Article 19 (35 U.S.C. 371(c)(3))</p> <p>a <input type="checkbox"/> are transmitted herewith (required only if not transmitted by the International Bureau)</p> <p>b <input type="checkbox"/> have been transmitted by the International Bureau.</p> <p>c <input type="checkbox"/> have not been made, however, the time limit for making such amendments has NOT expired.</p> <p>d <input checked="" type="checkbox"/> have not been made and will not be made.</p> <p>8 <input type="checkbox"/> A translation of the amendments to the claims under PCT Article 19 (35 U.S.C. 371(c)(3)).</p> <p>9 <input type="checkbox"/> An oath or declaration of the inventor(s) (35 U.S.C. 371(c)(4)).</p> <p>10 <input type="checkbox"/> A translation of the annexes to the International Preliminary Examination Report under PCT Article 36 (35 U.S.C. 371(c)(5)).</p> <p><b>Items 11. to 16. below concern document(s) or information included:</b></p> <p>11 <input type="checkbox"/> An Information Disclosure Statement under 37 CFR 1.97 and 1.98.</p> <p>12 <input type="checkbox"/> An assignment document for recording. A separate cover sheet in compliance with 37 CFR 3.28 and 3.31 is included</p> <p>13 <input checked="" type="checkbox"/> A <b>FIRST</b> preliminary amendment.</p> <p><input type="checkbox"/> A <b>SECOND</b> or <b>SUBSEQUENT</b> preliminary amendment.</p> <p>14 <input type="checkbox"/> A substitute specification</p> <p>15 <input type="checkbox"/> A change of power of attorney and/or address letter.</p> <p>16 <input checked="" type="checkbox"/> Other items or information.</p> <p>a. WO 99/40198-Front Page with Abstract, Specification, References, Claims and Drawings (100 pp.)</p> <p>b. International Search Report (7 pp.)</p> <p>c. International Preliminary Examination Report and Annex (17 pp.)</p>		

09/601490-092400

Form PTO-1390 (REV 10-95) page 2 of 2

IN THE UNITED STATES PATENT AND TRADEMARK OFFICE

In re application of :  
Désiré José COLLEN : IDENTIFICATION, PRODUCTION  
: AND USE OF STAPHYLOKINASE  
International Application : DERIVATIVES WITH REDUCED  
No. PCT/EP99/00748 : IMMUNOGENICITY AND/OR  
: REDUCED CLEARANCE  
International Filing Date :  
04 February 1999 :  
Priority Dates Claimed :  
04 February 1998 :  
06 February 1998 :  
Serial No. Not Yet Assigned :  
Filed Concurrently Herewith :

Pittsburgh, Pennsylvania  
August 3, 2000

LETTER RECOGNIZING ATTORNEYS

Box PCT  
Assistant Commissioner for Patents  
Washington DC 20231

Sir:


Enclosed are appropriate papers for initiating the national phase of the above-identified PCT application, comprising a specification, claims and drawings. A Preliminary Amendment is also enclosed.

Please accept the application for purposes of granting a filing date and recognize Barbara E. Johnson, Richard L. Byrne, Jesse A. Hirshman, Registration Nos. 31,198, 28,498 and 40,016, respectively, as attorneys in this application, pending the filing of a formal Declaration and Power of Attorney.

Kindly direct all communications relating to this application to Barbara E. Johnson.

Respectfully submitted,  
WEBB ZIESENHEIM LOGSDON  
ORKIN & HANSON, P.C.

By

  
Barbara E. Johnson, Reg. No. 31,198  
Attorney for Applicants  
700 Koppers Building  
436 Seventh Avenue  
Pittsburgh, PA 15219-1818  
Telephone: 412/471-8815  
Facsimile: 412/471-4094

007260 06470960

PATENT APPLICATION/PCT  
Attorney Docket No. 702-001463

IN THE UNITED STATES PATENT AND TRADEMARK OFFICE

In re application of :  
Désiré José COLLEN : IDENTIFICATION, PRODUCTION  
: AND USE OF STAPHYLOKINASE  
International Application : DERIVATIVES WITH REDUCED  
No. PCT/EP99/00748 : IMMUNOGENICITY AND/OR  
: REDUCED CLEARANCE  
International Filing Date :  
04 February 1999 :  
Priority Dates Claimed :  
04 February 1998 :  
06 February 1998 :  
Serial No. Not Yet Assigned :  
Filed Concurrently Herewith :

Pittsburgh, Pennsylvania

August 3, 2000

PRELIMINARY AMENDMENT

BOX PCT  
Assistant Commissioner for Patents  
Washington, DC 20231

Sir:

Prior to initial examination, please amend the above-identified patent application as follows:

IN THE CLAIMS:

Please cancel original claims 1-27. Amended claims 1-30 were submitted during Chapter II proceedings with a letter dated 01 May 2000. Please further amend amended claims 6, 14, 27 and 29 as follows:

Claim 6, line 2, delete "claims 1-5" and insert therefor --claim 1--.

Claim 14, line 2, delete "claims 9-13" and insert therefor --claim 9--.

Claim 27, line 2, delete "claims 1 to 8" and insert therefor --claim 1--.

Claim 29, line 3, delete "claims 1 to 25" and insert therefor --claim 1--.

IN THE ABSTRACT:

After the claims, please insert a page containing the Abstract Of The Disclosure, which is attached hereto as a separately typed page.

REMARKS

Original claims 1-27 have been canceled by this Preliminary Amendment. Amended claims 6, 14, 27 and 29 have been further amended to remove the multiple dependencies for filing purposes. The Examiner and the Application Branch are respectfully requested not to enter the original claims 1-27.

A further Preliminary Amendment will be submitted in due course to address the amino acid sequence listings.


An Abstract Of The Disclosure has been added as a separately typed page to be inserted after the claims.

Entry of this Preliminary Amendment is respectfully requested.

Respectfully submitted,

WEBB ZIESENHEIM LOGSDON  
ORKIN & HANSON, P.C.

By

  
Barbara E. Johnson, Reg. No. 31,198  
Attorney for Applicant  
700 Koppers Building  
436 Seventh Avenue  
Pittsburgh, PA 15219-1818  
Telephone: 412/471-8815  
Facsimile: 412/471-4094

IDENTIFICATION, PRODUCTION AND USE OF STAPHYLOKINASE DERIVATIVES WITH REDUCED IMMUNO-GENICITY AND/OR REDUCED CLEARANCE

5           The present invention relates to new  
staphylokinase derivatives with reduced immunogenicity  
which can be administered by continuous infusion or by  
single intravenous bolus injection, to their  
identification, production and use in the treatment of  
10 arterial thrombosis and to the preparation of a  
pharmaceutical composition for treating arterial  
thrombosis. More in particular the invention relates to  
the use of engineered staphylokinase derivatives for the  
preparation of a pharmaceutical composition for treating  
15 myocardial infarction.

          Staphylokinase, a protein produced by certain  
strains of Staphylococcus aureus, which was shown to have  
profibrinolytic properties more than 4 decades ago (1, 2)  
appears to constitute a potent thrombolytic agent in  
20 patients with acute myocardial infarction (3, 4). The  
staphylokinase gene has been cloned from the  
bacteriophages sak $\phi$ C (5) and sak42D (6) as well as from  
the genomic DNA (sakSTAR) of a lysogenic Staphylococcus  
aureus strain (7). The staphylokinase gene encodes a  
25 protein of 163 amino acids, with amino acid 28  
corresponding to the NH<sub>2</sub>-terminal residue of full length  
mature staphylokinase (6, 8, 9). The mature protein  
sequence of the wild-type variant SakSTAR (9) is  
represented in Figure 1. Only four nucleotide differences  
30 were found in the coding regions of the sak $\phi$ C, sak42D and  
sakSTAR genes, one of which constituted a silent mutation  
(6, 8, 9). In a plasma milieu, staphylokinase is able to  
dissolve fibrin clots without associated fibrinogen  
degradation (10-12). This fibrin-specificity of  
35 staphylokinase is the result of reduced inhibition by  
 $\alpha_2$ -antiplasmin of plasmin.staphylokinase complex bound to  
fibrin, recycling of staphylokinase from the  
plasmin.staphylokinase complex following inhibition by

007250 06470960

007260 0640960

$\alpha_2$ -antiplasmin, and prevention of the conversion of circulating plasminogen.staphylokinase to plasmin.staphylokinase by  $\alpha_2$ -antiplasmin (13-15). In addition staphylokinase has a weak affinity for circulating but a high affinity for fibrin-bound plasminogen (16) and staphylokinase requires  $\text{NH}_2$ -terminal processing by plasmin to display its plasminogen activating potential (17). In several experimental animal models, staphylokinase appears to be equipotent to streptokinase for the dissolution of whole blood or plasma clots, but significantly more potent for the dissolution of platelet-rich or retracted thrombi (18, 19). Staphylokinase is a heterologous protein and is immunogenic in man. The intrinsic immunogenicity of staphylokinase, like that of streptokinase, clearly hampers its unrestricted use. Not only will patients with preexisting high antibody titers be refractory to the thrombolytic effect of these agents, but allergic side effects and occasional life-threatening anaphylaxis may occur (20). Because both streptokinase and staphylokinase are heterologous proteins, it is not obvious that their immunogenicity could be reduced by protein engineering. Indeed, no successful attempts to generate active low molecular weight fragments from streptokinase have been reported. In staphylokinase, deletion of the  $\text{NH}_2$ -terminal 17 amino acids or the  $\text{COOH}$ -terminal 2 amino acids inactivates the molecule, which in addition is very sensitive to inactivation by site-specific mutagenesis (21).

It is therefore the object of the present invention to provide less immunogenic variants of staphylokinase having preferably a higher specific activity and/or lower plasma clearance and/or increased thrombolytic potency.

In the research that ultimately led to the present invention it was already found that the wild-type staphylokinase variant SakSTAR (9) contains three non-overlapping immunodominant epitopes, at least two of

which can be eliminated by specific site-directed mutagenesis, without inactivation of the molecule. This has been disclosed in EP-95200023.0 (22). These engineered staphylokinase variants are less reactive with antibodies elicited in patients treated with wild-type staphylokinase, and are significantly less immunogenic than wild-type staphylokinase, as demonstrated in rabbit and baboon models and in patients with peripheral arterial occlusion (22).

The present invention now relates to general methods for the identification, production and use of staphylokinase derivatives showing a reduced antigenicity and immunogenicity as compared to wild-type staphylokinase as well as for variants with selective derivatization with polyethylene glycol. The derivatives preferably have a higher specific activity and/or lower plasma clearance and/or increased thrombolytic potency. The derivatives have essentially the amino acid sequence of wild-type staphylokinase or modified versions thereof and essentially intact biological activities, but have a reduced reactivity with a panel of murine monoclonal antibodies and/or with antibodies induced in patients by treatment with wild-type SakSTAR. The polyethylene glycol substituted ("pegylated") variants have reduced plasma clearances rendering them particularly suited for use by single intravenous bolus administration. Instead of PEG other pharmaceutically acceptable macromolecules can be used.

More in particular, the invention provides for staphylokinase derivatives SakSTAR(K35X,G36X,E65X,K74X,E80X,D82X,K102X,E108X,K109X,K121X,K130X, K135X,K136X,+137X) having the amino acid sequence as depicted in figure 1 in which the amino acids Lys in position 35, Gly in position 36, Glu in position 65, Lys in position 74, Glu in position 80, Asp in position 82, Lys in position 102, Glu in position 108, Lys in position 109, Lys in position 121, Lys in position 130, Lys in position 135 and/or Lys in position 136 have been replaced with other amino acids and/or in which one amino acid has been added at the COOH-terminus, thus altering the immunogenicity

001260" 06470560



after administration in patients, without markedly reducing the specific activity.

Further preferred embodiments of the invention are staphylokinase derivatives listed in Tables 1, 3, 4, 5, 6, 7, 8, 13, 19 and 20, having the amino acid sequence as depicted in figure 1 in which the indicated amino acids have been replaced by other amino acids thus reducing the absorption of SakSTAR-specific antibodies from plasma of patients treated with staphylokinase, without reducing the specific activity.

Derivatives in which the specific activity is increased and the immunogenicity is decreased are the following:

- SakSTAR(K74A,E75A,R77A), SakSTAR(K35A,E75A),  
 15 SakSTAR(E75A), SakSTAR(E80A,D82A), SakSTAR(E80A),  
 SakSTAR(D82A), SakSTAR(E75A,D82A), SakSTAR(S34G,G36R,  
 H43R), SakSTAR(K35A), SakSTAR(E80A), SakSTAR(D82A,S84A),  
 SakSTAR(T90A), SakSTAR(Y92A), SakSTAR(K130A),  
 SakSTAR(V132A), SakSTAR(S34G,G36R,H43R), SakSTAR(G36R),  
 20 SakSTAR(H43R), SakSTAR(G36R,K74R), SakSTAR(K35E),  
 SakSTAR(K74Q), SakSTAR(K130T), SakSTAR(V132L),  
 SakSTAR(V132T), SakSTAR(V132N), SakSTAR(V132R),  
 SakSTAR(K130T,K135R), SakSTAR(G36R,K130T,K135R),  
 SakSTAR(K74R,K130T,K135R), SakSTAR(K74Q,K130T,K135R),  
 25 SakSTAR(G36R,K74R,K130T,  
 K135R), SakSTAR(G36R,K74Q,K130T,K135R), SakSTAR(G36R,  
 H43R,K74R,K130T,K135R), SakSTAR(E65A,K74Q,K130T,K135R),  
 SakSTAR(E65Q,K74Q,K130T,K135R), SakSTAR(K74Q,K86A,  
 K130T,K135R), SakSTAR(E65Q,T71S,K74Q,K130T,K135R),  
 30 SakSTAR(K74Q,K130A,K135R), SakSTAR(E65Q,K74Q,K130A,  
 K135R), SakSTAR(K74Q,K130E,K135R), SakSTAR(K74Q,K130E,  
 V132R,K135R), SakSTAR(E65Q,K74Q,T90A,K130A,K135R),  
 SakSTAR(E65Q,K74Q,N95A,K130A,K135R), SakSTAR(E65Q,K74Q,  
 E118A,K130A,K135R), SakSTAR(E65Q,K74Q,N95A,E118A,K130A,  
 35 K135R), SakSTAR(N95A,K130A,K135R), SakSTAR(E65Q,K74Q,  
 K109A,K130,K135R), SakSTAR(E65Q,K74Q,E108A,K109A,

- K130T, K135R), SakSTAR(E65Q, K74Q, K121A, K130T, K135R),  
 SakSTAR(E65Q, K74Q, N95A, E118A, K130A, K135R, K136A, +137K),  
 SakSTAR(E80A, D82A, K130T, K135R), SakSTAR(K74R, E80A, D82A,  
 K130T, K135R), SakSTAR(K74Q, E80A, D82A, K130T, K135R),  
 5 SakSTAR(K35A, K74R, E80A, D82A, K130T, K135R), SakSTAR(E65D,  
 K74R, E80A, D82A, K130T, K135R), SakSTAR(E65S, K74R, E80A,  
 D82A, K130T, K135R), SakSTAR(S34G, G36R, K74R, K130T, K135R),  
 SakSTAR(E65A, K74R, E80A, D82A, K130T, K135R), SakSTAR(E65N,  
 K74R, E80A, D82A, K130T, K135R), SakSTAR(E65Q, K74R, E80A,  
 10 D82A, K130T, K135R), SakSTAR(K57A, E58A, E61A, E80A, D82A,  
 K130T, K135R), SakSTAR(E65D, K74Q, E80A, D82A, K130T, K135R),  
 SakSTAR(E65Q, K74Q, E80A, D82A, K130T, K135R), SakSTAR(K35A,  
 E65D, K74Q, E80A, D82A, K130T, K135R), SakSTAR(K74R, E80A, D82A,  
 S103A, K130T, K135R), SakSTAR(E65D, K74R, E80A, D82A, K109A,  
 15 K130T, K135R), SakSTAR(E65D, K74R, E80A, D82A, K130T,  
 K135R, K136A), SakSTAR(E65Q, K74Q, D82A, S84A, K130T, K135R),  
 SakSTAR(K35A, K74Q, E80A, D82A, K130T, K135R), SakSTAR(K35A,  
 E65D, K74R, E80A, D82A, K130T, K135R).

- Of these SakSTAR(E65D, K74R, E80A, D82A, K130T,  
 20 K135R) having the code SY19 and SakSTAR(K35A, E65Q, K74R,  
 E80A, D82A, T90A, E99D, T101S, E108A, K109A, K130T, K135R) having  
 the code SY161 are especially preferred.

- Besides the above described substitution  
 derivatives the invention relates to derivatives having  
 25 in addition an amino acid substituted with Cys. This type  
 of substitution may result in dimerization and/or  
 increased specific activity and/or reduced clearance  
 and/or increased thrombolytic potency. Reduced plasma  
 clearance is in particular obtained when the derivative  
 30 is substituted with polyethylene glycol.

- Preferred embodiments of such staphylokinase  
 derivatives are those wherein the Cys is chemically  
 modified with polyethylene glycol with molecular weights  
 up to 20 kDa. In particular embodiments selected amino  
 35 acids in the NH<sub>2</sub>-terminal region of 10 amino acids, are  
 substituted with Cys, which is chemically modified with  
 polyethylene glycol with molecular weights up to 20 kDa.  
 These derivatives are characterized by a significantly

reduced plasma clearance and maintained thrombolytic potency upon single intravenous bolus administration at a reduced dose.

More in particular the serine in position 2 or  
 5 3 is substituted with a cysteine and the cysteine is chemically modified with polyethylene glycol having a molecular weight of 5, 10 or 20 kDa. Preferred embodiments of these derivatives are SY161(S3C-MP5), SY161(S3C-P10), SY161(S3C-P20), SY19(S3C-MP5), SY19(S3C-  
 10 SP5), SY19(S2C-SP5, S3C-SP5), SY19(S3C-P20), SY19(S3C-P10) all as defined in table 20.

The presence of cysteines allows the formation of dimers of two staphylokinase derivatives of the invention.

15 The invention also relates to a method for producing the derivatives of the invention by preparing a DNA fragment comprising at least the part of the coding sequence of staphylokinase that provides for its biological activity; performing in vitro site-directed  
 20 mutagenesis on the DNA fragment to replace one or more codons for wild-type amino acids by a codon for another amino acid; cloning the mutated DNA fragment in a suitable vector; transforming or transfecting a suitable host cell with the vector; culturing the host cell under  
 25 conditions suitable for expressing the DNA fragment; purifying the expressed staphylokinase derivative to homogeneity and derivatizing the variant with polyethylene glycol.

Preferably the DNA fragment is a 453 bp  
 30 EcoRI-HindIII fragment of the plasmid pMEX602sakB (22, 23), the in vitro site-directed mutagenesis is preferably performed by spliced overlap extension polymerase chain reaction. Such overlap extension PCR is preferably performed with Vent DNA polymerase (New England Biolabs)  
 35 or Taq polymerase (Boehringer Mannheim) and with available or generated wildtype sakSTAR or sakSTAR variants as template (24).

007260 0640960

The invention also relates to pharmaceutical compositions comprising at least one of the staphylokinase derivatives according to the invention together with a suitable excipient, for treatment of

5 arterial thrombosis. Pharmaceutical compositions, containing less immunogenic staphylokinase variants or "pegylated" staphylokinase variants as the active ingredient, for treating arterial thrombosis in human or veterinary practice may take the form of powders or

10 solutions and may be used for intravenous, intraarterial or parenteral administration. Such compositions may be prepared by combining (e.g. mixing, dissolving etc.) the active compound with pharmaceutically acceptable excipients of neutral character (such as aqueous or

15 non-aqueous solvents, stabilizers, emulsifiers, detergents, additives), and further, if necessary with dyes.

Furthermore the invention relates to the use of the staphylokinase derivatives for the treatment of

20 arterial thrombosis, in particular myocardial infarction, and to the use of staphylokinase derivatives for the preparation of a pharmaceutical composition for the treatment of arterial thrombosis, in particular myocardial infarction. In the above and the following the

25 terms "derivatives", "mutants" and "variants" are used interchangeably.

Based on the present invention other variants and improvements will be obvious for the person skilled in the art. Thus random mutagenesis is likely to generate

30 alternative mutants with reduced immunogenicity and possibly increased functional activity, whereas deletions or substitution with other amino acids may yield additional variants with reduced immunogenicity.

The present invention will be demonstrated in

35 more detail in the following examples, that are however not intended to be limiting to the scope of the invention. In the Examples reference is made to the following figures:

Fig 1. Protein sequence of wild-type staphylokinase, SakSTAR. Numbering starts with the NH<sub>2</sub>-terminal amino acid of mature full length staphylokinase.

5 Fig 2. Time course of neutralizing activities (left panel) and specific IgG against administered agent (right panel) following intra-arterial infusion of SakSTAR (open circles, n= 9), SakSTAR(K74A) (closed circles, n= 11) or SakSTAR(K74A,E75A, R77A) (open  
10 squares, n= 6) in patients with peripheral arterial occlusion. The data represent median values and interquartile ranges, in  $\mu\text{g/mL}$ .

Fig 3. Protein sequence of wild-type staphylokinase, SakSTAR with indicated amino acid  
15 substitutions.

squares: single amino acid substitutions;  
circles: combined (2 to 3) amino acid to Ala substitutions.

Fig. 4. Temperature stability of SakSTAR, (A);  
20 SakSTAR(K74Q,E80A,D82A,K130T, K135R) (B);  
SakSTAR(E65D,K74R,E80A,D82A,K130T,K135R), (C); and  
SakSTAR(K35A,E65D,K74Q,E80A,D82A, K130T,K135R), (D).  
(○): 4°C; (●): 20°C; (▽): 37°C; (▼): 56°C; (□): 70°C.

Fig 5. Time course of neutralizing activities  
25 (left panel) and specific IgG against administered agent (right panel) following intra-arterial infusion of  
SakSTAR (circles, n= 6),  
SakSTAR(K74Q,E80A,D82A,K130T,K135R) (squares, n= 6) or  
SakSTAR(E65D,K74R,E80A,D82A,K130T,K135R) (triangles, n=  
30 6) in patients with peripheral arterial occlusion. The data represent median values and 15-85 percentile ranges, in  $\mu\text{g/mL}$ .

### EXAMPLES

#### 35 EXAMPLE 1

##### Epitope mapping of wild-type staphylokinase

The epitope specificity of a panel of 15 murine MAbs (22) raised against wild-type SakSTAR was determined

004260" 064T0960

by real-time biospecific interaction analysis (BIA) with the BIAcore instrument (Pharmacia, Biosensor AB, Uppsala, Sweden). The MAbs were immobilized on the surface of the Sencor Chip CM5 with the Amine Coupling Kit (Pharmacia Biosensor AB) as recommended by the manufacturer (25). Immobilization was performed from protein solutions at a concentration of 20  $\mu\text{g/mL}$  in 10 mmol/L sodium acetate at pH 5.0 at a flow rate of 5  $\mu\text{L/min}$  during 6 minutes. This resulted in covalent attachment of 5,000 to 10,000 resonance unit (RU) of antibody (corresponding to 0.035 to 0.07 pmol/ $\text{mm}^2$ ). The SakSTAR solutions were passed by continuous flow at 20°C past the sensor surface. At least four concentrations of each analyte (range, 50 nmol/L to 50 mol/L) in 10 mmol/L HEPES, 3.4 mmol/L EDTA, 0.15 mol/L NaCl, and 0.005% Surfactant P20, pH 7.2, were injected at a flow rate of 5  $\mu\text{L/min}$  during 6 minutes in the association phase. Then sample was replaced by buffer, also at a flow rate of 5  $\mu\text{L/min}$  during 6 minutes. After each cycle, the surface of the sensor chip was regenerated by injection of 5  $\mu\text{L}$  of 15 mmol/L HCl. Apparent association ( $k_{\text{ass}}$ ) and apparent dissociation ( $k_{\text{diss}}$ ) rate constants were derived from the sensorgrams as described in detail elsewhere (26), and association equilibrium constants ( $K_A$ ) calculated as their ratio.

Determination of the equilibrium association constants for the binding of wild-type and variant SakSTAR to insolubilized MAbs (Table 1) yielded apparent association constants of  $10^7$  to  $10^8$  (mol/L) $^{-1}$ , which are one to two orders of magnitude lower than the apparent association constants previously obtained for the binding of these MAbs to insolubilized wild-type SakSTAR (22). If the MAbs instead of the SakSTAR variants are insolubilized, avidity effects of the bivalent MAbs are avoided. The present values are indeed in better agreement with known association constants of MAbs, and therefore this "reversed" procedure was used throughout the present invention.

In the tables the column indicated with "Variant" states the various staphylokinase derivatives which are identified by listing between brackets the substituted amino acids in single letter symbols followed by their position number in the mature staphylokinase sequence and by the substituting amino acids in single letter symbol; the column "Exp." indicates expression levels in mg/L, and the column "Spec. Act." indicates the specific activity in Home Units as defined in example 2.

Indications "17G11", "26A2" etc. refer to monoclonal antibodies binding to the indicated epitopes I, II and III as defined in reference 22. Epitope I is recognized by the antibody cluster 17G11, 26A2, 30A2, 2B12 and 3G10, whereas epitope II is recognized by the antibody cluster 18F12, 14H5, 28H4, 32B2 and 7F10, and epitope III by the antibody cluster 7H11, 25E1, 40C8, 24C4 and 1A10. Human plasma "Pool" indicates a plasma pool from initially 16 and subsequently 10 patients immunized by treatment with SakSTAR, "Subpool B" indicates a plasma pool from three patients that absorbed less than 50% of the induced antibodies with SakSTAR(K35A,E38A,K74A,E75A,R77A) and "Subpool C" indicates a plasma pool from 3 patients that absorbed >90% of the induced antibodies with SakSTAR(K35A,E38A,K74A,E75A,R77A) (22).

In tables 6, 7 and 8 an additional pool of plasma from 40 patients immunized by treatment with SakSTAR (Pool 40) was also used.

## EXAMPLE 2

Construction, epitope mapping with murine monoclonal antibodies and absorption with pooled plasma of immunized patients, of "alanine-to-wild-type" reversal variants of "charged-cluster-to-alanine" mutants of staphylokinase

### 1. Introduction

As stated above, wild-type staphylokinase (SakSTAR variant (9)) contains three non-overlapping immunodominant epitopes, two of which can be eliminated by specific site-directed substitution of clusters of two

007250" 05470550

(K35A,E38A or E80A,D82A) or three (K74A,E75A,R77A) charged amino acids with Ala (22). The combination mutants SakSTAR(K35A,E38A,K74A,E75A,R77A) in which Lys35, Glu38, Lys74, Glu75 and Arg77, and SakSTAR(K74A,E75A, R77A,E80A,D82A) in which Lys74, Glu75, Arg77, Glu80 and Asp82 were substituted with Ala (previously identified as SakSTAR.M3.8 and SakSTAR.M8.9, respectively (22)), were found to have a reduced reactivity with murine monoclonal antibodies against two of the three immunodominant epitopes and to absorb on average only 2/3 of the neutralizing antibodies elicited in 16 patients by treatment with wild-type SakSTAR (22). These mutants also induced less antibody formation than wild-type SakSTAR in experimental thrombolysis models in rabbits and baboons, and in patients with peripheral arterial occlusion (22). However, their specific activities were reduced to approximately 50% of that of wild-type SakSTAR, which would be of some concern with respect to the clinical use of these compounds.

In an effort to improve the activity and stability without loss of the reduced antibody recognition, the effect of a systematic reversal of one or more of these substituted amino acids to the wild-type residues was studied. Fourteen new mutants were constructed, purified and characterized in terms of specific activity, reactivity with the panel of murine monoclonal antibodies, and absorption of antibodies from plasma of patients treated with wild-type SakSTAR (Table 1). The present example thus focusses on reversal from alanine to the wild-type residue of one or more of the seven amino acids of SakSTAR listed above i.e. K35, E38, K74, E75, R77, E80 and D82.

## 2. Reagents and Methods

The source of all reagents used in the present study has previously been reported (22). Restriction enzymes were purchased from Pharmacia (Uppsala, Sweden) or Boehringer Mannheim (Mannheim, Germany). T4 DNA



ligase, Klenow Fragment of E. coli DNA polymerase I and alkaline phosphatase were obtained from Boehringer Mannheim. Enzyme reactions were performed using the conditions suggested by the suppliers. Plasmid DNA was isolated using a QIAGEN-purification protocol (provided by Westburg, Leusden, The Netherlands). pMEX.602sakB (i.e. pMEX.SakSTAR) was constructed as described elsewhere (23). SakSTAR, SakSTAR(K35A,E38A), SakSTAR(K74A,E75A,R77A), SakSTAR(E80A,D82A), SakSTAR(K35A,E38A,K74A,E75A,R77A) and SakSTAR(K74A,E75A,R77A,E80A,D82A) were produced and purified as described elsewhere (22). Transformations of *E. coli* were performed utilizing the calcium phosphate procedure. DNA sequencing was performed using the dideoxy chain termination reaction method and the Automated Laser fluorescent A.L.F.<sup>TM</sup> (Pharmacia). The chromogenic substrate (S2403) L-Pyroglutamyl-L-phenylalanyl-L-lysine-p-nitroaniline hydrochloride was purchased from Chromogenix (Belgium). <sup>125</sup>I-labeled fibrinogen was purchased from Amersham (UK). All other methods used in the present example have been previously described (22,27).

### 3. Construction of expression plasmids

The plasmids encoding SakSTAR(K35A,E38A,K74A,E75A), SakSTAR(E38A,E75A,R77A), SakSTAR(E38A,E75A), SakSTAR(K35A,E75A,R77A), SakSTAR(K35A,E75A), SakSTAR(E80A), SakSTAR(D82A), SakSTAR(E75A,D82A), SakSTAR(K74A) and SakSTAR(E75A) were constructed by the spliced overlap extension polymerase chain reaction (SOE-PCR) (24), using Vent DNA polymerase (New England Biolabs, Leusden, The Netherlands), and available or generated sakSTAR variants as template. Two fragments were amplified by PCR, the first one starting from the 5' end of the staphylokinase gene with primer 5'-CAGGAAACAGAATTCAGGAG-3' to the region to be mutagenized (forward primer), the second one from the same region (backward primer) to the 3' end of the

staphylokinase gene with primer  
 5'-CAAAACAGCCAAGCTTCATTCATTCAGC-3'. The forward and  
 backward primers shared an overlap of around 24 bp (pri-  
 mers not shown). The two purified fragments were then  
 5 assembled together in a new primerless PCR using Taq  
 polymerase (Boehringer Mannheim). After 7 cycles (1 min  
 at 94°C, 1 min at 70°C), the extended product was  
 reamplified by adding the 5' and 3' end primers (see  
 above) to the PCR reaction and by cycling 25 times (1 min  
 10 at 94°C, 1 min 55°C, 1 min at 72°C). The final product  
 was purified, digested with EcoRI and HindIII and cloned  
 into the corresponding sites of pMEX602sakB. The plasmid  
 encoding SakSTAR(E38A,K74A,E75A,R77A) was assembled by  
 digestion of pMEX602sakB and pMEX.SakSTAR(K35A,E38A,  
 15 K74A,E75A,R77A) with BpmI which cuts between the codons  
 for K35 and E38 of SakSTAR, and ligation of the required  
 fragments. The plasmid encoding SakSTAR(K35A,K74A,E75A,  
 R77A) was constructed by digestion of pMEX.SakSTAR(K35A,  
 E38A,K74A,E75A,R77A) and pMEX.SakSTAR(K74A,E75A,R77A)  
 20 with BpmI and religation of the required fragments. The  
 plasmids encoding SakSTAR(K35A,E38A, E75A,R77A) and  
 SakSTAR(K35A,E38A,K74A,R77A) were constructed by two PCR  
 using pMEX.SakSTAR(K35A,E38A,K74A,E75A,R77A) as template,  
 followed by restriction ligation and recloning into  
 25 pMEX602sakB.

#### 4. Expression and purification of SakSTAR variants

The SakSTAR variants were expressed and  
 purified, as described below, from transformed *E. coli*  
 30 WK6 grown either in LB medium [SakSTAR(E38A,K74A,E75A,  
 R77A), SakSTAR(K74A), SakSTAR(E75A) and SakSTAR(E75A,  
 D82A)], or in terrific broth (TB) (28) medium  
 [SakSTAR(K35A,K74A,E75A,R77A), SakSTAR(K35A,E38A,E75A,  
 R77A), SakSTAR(K35A,E38A,K74A,R77A), SakSTAR(K35A,  
 35 E38A,E75A), SakSTAR(E38A,E75A, R77A), SakSTAR(E38A,E75A),  
 SakSTAR(K35A,E75A,R77A), SakSTAR(K35A,E75A),  
 SakSTAR(E80A), and SakSTAR(D82A)].

For the derivatives produced in TB medium, a 4  
15 mL aliquot of an overnight saturated culture in LB medium  
was used to inoculate a 2 L culture in terrific broth  
containing 100  $\mu$ g/mL ampicillin. The culture was grown  
with vigorous aeration for 20 hours at 30°C. The cells  
were pelleted by centrifugation, resuspended in 1/10  
20 volume (200 mL) of 0.01 mol/L phosphate buffer pH 6.5 and  
disrupted by sonication at 0°C. The suspension was then  
centrifuged for 20 min at 20,000 rpm and the supernatant  
was stored at -20°C until purification. Cleared cell  
lysates containing the SakSTAR variants were subjected to  
25 chromatography on a 1.6 x 6 cm column of SP-Sephadex,  
followed by chromatography on a 1.6 x 5 cm column of  
Q-Sepharose [variants SakSTAR(E38A,K74A,E75A,R77A),  
SakSTAR(K35A,K74A,E75A, R77A), SakSTAR(K35A,E38A,  
E75A,R77A), SakSTAR(K35A,E38A,K74A,R77A) and  
30 SakSTAR(K35A, E38A,K74A,E75A)] or by chromatography on a  
1.6 x 6 cm column of phenyl-Sepharose [variants Sak-  
STAR(E35A,E38A,R77A), SakSTAR(E38A,E75A), SakSTAR-  
(K35A,E75A,R77A), SakSTAR(K35A,E75A), SakSTAR(K74A),  
SakSTAR(E75A), SakSTAR(E80A), SakSTAR(D82A) and Sak-  
35 STAR(E75A,D82A)]. The SakSTAR containing fractions,  
localized by SDS-gel electrophoresis, were pooled for  
further analysis.

For the derivatives produced in TB medium, a 4  
15 mL aliquot of an overnight saturated culture in LB medium  
was used to inoculate a 2 L culture in terrific broth  
containing 100  $\mu$ g/mL ampicillin. The culture was grown  
with vigorous aeration for 20 hours at 30°C. The cells  
were pelleted by centrifugation, resuspended in 1/10  
20 volume (200 mL) of 0.01 mol/L phosphate buffer pH 6.5 and  
disrupted by sonication at 0°C. The suspension was then  
centrifuged for 20 min at 20,000 rpm and the supernatant  
was stored at -20°C until purification. Cleared cell  
lysates containing the SakSTAR variants were subjected to  
25 chromatography on a 1.6 x 6 cm column of SP-Sephadex,  
followed by chromatography on a 1.6 x 5 cm column of  
Q-Sepharose [variants SakSTAR(E38A,K74A,E75A,R77A),  
SakSTAR(K35A,K74A,E75A, R77A), SakSTAR(K35A,E38A,  
E75A,R77A), SakSTAR(K35A,E38A,K74A,R77A) and  
30 SakSTAR(K35A, E38A,K74A,E75A)] or by chromatography on a  
1.6 x 6 cm column of phenyl-Sepharose [variants Sak-  
STAR(E35A,E38A,R77A), SakSTAR(E38A,E75A), SakSTAR-  
(K35A,E75A,R77A), SakSTAR(K35A,E75A), SakSTAR(K74A),  
SakSTAR(E75A), SakSTAR(E80A), SakSTAR(D82A) and Sak-  
35 STAR(E75A,D82A)]. The SakSTAR containing fractions,  
localized by SDS-gel electrophoresis, were pooled for  
further analysis.

## 5. Physicochemical and biochemical analysis

Protein concentrations were determined according to Bradford (29). The specific activities of SakSTAR solutions were determined with a chromogenic substrate assay carried out in microtiter plates using a mixture of 80  $\mu$ L SakSTAR solution and 100  $\mu$ L Glu-plasminogen solution prepared as described elsewhere (30) (final concentration 0.5  $\mu$ mol/L). After incubation for 30 min at 37°C, generated plasmin was quantitated by addition of 20  $\mu$ L S2403 (final concentration 1 mmol/L) and measurement of the absorption at 405 nm. The activity was expressed in home units (HU) by comparison with an in-house standard (lot STAN5) which was assigned an activity of 100,000 HU (100 KHU) per mg protein as determined by amino acid composition (7). SDS-PAGE was performed with the Phast System™ (Pharmacia, Uppsala, Sweden) using 10-15% gradient gels and Coomassie Brilliant blue staining. Reduction of the samples was performed by heating at 100°C for 3 min in the presence of 1% SDS and 1% dithiothreitol. The specific activities of the different SakSTAR mutants determined with the chromogenic substrate assay are summarized in Table 1.

## 6. Binding to murine monoclonal antibodies

In agreement with previous observations (22), SakSTAR(K74A,E75A,R77A) did not react with 4 of the 5 MAb's recognizing epitope I, whereas SakSTAR(K35A,E38A) did not react with 3 of the 5 and SakSTAR(E80A,D82A) not with 4 of the 5 MAb's recognizing epitope III. These reduced reactivities were additive in SakSTAR(K35A,E38A,K74A,E75A,R77A) and in SakSTAR(K74A,E75A,R77A,E80A,D82A). The reduced reactivity of SakSTAR(K74A,E75A,R77A) was fully maintained in SakSTAR(K35A,E38A,K74A,E75A) and in SakSTAR(K35A,E75A,R77A), largely in SakSTAR(K35A,E38A,E75A,R77A), SakSTAR(E38A,E75A,R77A), SakSTAR(E38A,E75A) and SakSTAR(E75A), but much less in SakSTAR(K35A,E38A,K74A,R77A) and SakSTAR(K74A), indicating that E75 is the main contributor to the binding of the 4 MAb's recognizing

epitope I of SakSTAR. However, surprisingly, binding of epitope I antibodies to SakSTAR(E75A,D82A) was normal in two independent preparations from expression plasmids with confirmed DNA sequences. The reduced reactivity of the 3 MAbs of epitope III with SakSTAR(K35A,E38A) required both K35 and E38, as demonstrated with SakSTAR(E38A,K74A,E75A,R77A) and SakSTAR(K35A,K74A,E75A,R77A), with SakSTAR(E38A,E75A) and SakSTAR(K35A,E75A) and with SakSTAR(E38A,E75A,R77A) and SakSTAR(K35A,E75A,R77A). The reduced reactivity of the 4 MAbs of cluster III with SakSTAR(E80A,D82A) was maintained in SakSTAR(D82A) but not in SakSTAR(E80A).

7. Absorption of antibodies, elicited in patients by treatment with wild-type SakSTAR

Plasma samples from 16 patients with acute myocardial infarction, obtained several weeks after treatment with SakSTAR (4, 31) were used. The staphylokinase-neutralizing activity in these samples was determined as follows. Increasing concentrations of wild-type or variant SakSTAR (50  $\mu$ L volumes containing 0.2 to 1000  $\mu$ g/mL) were added to a mixture of 300  $\mu$ L citrated human plasma and 50  $\mu$ L buffer or test plasma, immediately followed by addition of 100  $\mu$ L of a mixture containing thrombin (50 NIH units/mL) and  $\text{CaCl}_2$  (25 mmol/L). The plasma clot lysis time was measured and plotted against the concentration of SakSTAR moiety. From this curve the concentration of staphylokinase moiety that produced complete clot lysis in 20 min was determined. The neutralizing activity titer was determined as the difference between the test plasma and buffer values and was expressed in  $\mu$ g per mL test plasma. The results of the individual patients have been reported elsewhere (22). For the present invention, three plasma pools were made, one from 10 patients from whom sufficient residual plasma was available, one from three patients that absorbed less than 50% of the antibodies with SakSTAR(K35A,E38A, K74A,E75A,R77A) (Subpool B) and

001250 06410960

one from three patients that absorbed >90% of the antibodies with SakSTAR(K35A,E38A,K74A,E75A, R77A) (Subpool C). These plasma pools were diluted (1/30 to 1/200) until their binding to SakSTAR substituted chips in the BIAcore instrument amounted to approximately 2000 RU. From this dilution a calibration curve for antibody binding was constructed using further serial two-fold dilutions. The plasma pools were absorbed for 10 min with 100 nmol/L of the SakSTAR variants, and residual binding to immobilized SakSTAR was determined. Residual binding was expressed in percent of unabsorbed plasma, using the calibration curve.

The results are summarized in Table 1. Whereas wild-type SakSTAR absorbed more than 95% of the binding antibodies from pooled plasma of the 10 patients, incomplete absorption (<60%) was observed with SakSTAR(K74A,E75A,R77A), SakSTAR(K35A,E38A,K74A,E75A,R77A), SakSTAR(E38A,K74A,E75A,R77A), SakSTAR(K35A,K74A,E75A,R77A), SakSTAR(K35A,E38A,K74A,R77A), SakSTAR(K35A,E38A,K74A,E75A), SakSTAR(K74A) and SakSTAR(K74A,E75A,R77A,E80A,D82A) but absorption was nearly complete with SakSTAR(K35A,E38A), SakSTAR(K35A,E38A,E75A,R77A), SakSTAR(E38A,E75A,R77A), SakSTAR(E38A,E75A), SakSTAR(K35A,E75A,R77A), SakSTAR(K35A,E75A), SakSTAR(E75A), SakSTAR(E80A,D82A), SakSTAR(E80A), SakSTAR(D82A) and SakSTAR(E75A,D82A). These results, surprisingly, demonstrate that approximately 40% of the antibodies elicited in patients by treatment with wild-type SakSTAR depend on K74 for their binding (Table 1). Absorption with pooled plasma from 3 patients from which <50% of the antibodies were absorbed with SakSTAR(K35A,E38A,K74A,E75A,R77A) (Subpool B) confirmed the predominant role of K74 for antibody recognition. As expected, absorption with pooled plasma from 3 patients from which >95% of the antibodies were absorbed with SakSTAR(K35A,E38A,K74A,E75A,R77A) (Subpool C) was nearly complete with all variants tested.

004260" 0640960

**EXAMPLE 3****Comparative thrombolytic efficacy and immunogenicity of SakSTAR(K74A,E75A,R77A) and SakSTAR(K74A) versus SakSTAR in patients with peripheral arterial occlusion****5 1. Purification of SakSTAR(K74A,E75A,R77A) and SakSTAR(K74A) for use in vivo**

A 12 to 24 L culture (in 2 L batches) of the variants SakSTAR(K74A,E75A,R77A), or of SakSTAR(K74A) was grown and IPTG-induced in LB medium supplemented with 100  
10 µg/mL ampicillin, pelleted, resuspended, disrupted by sonication and cleared as described above. The compounds were purified by chromatography on a 5 x 20 cm column of SP-Sephadex, a 5 x 10 cm column of Q-Sepharose and/or a 5 x 13 cm column of phenyl-Sepharose using buffer systems  
15 described elsewhere (22, 23). The materials were then gel filtered on sterilized Superdex 75 to further reduce their endotoxin content. The SakSTAR variant containing fractions were pooled, the protein concentration was adjusted to 1 mg/mL and the material sterilized by  
20 filtration through a 0.22 µm Millipore filter. The methodology used to determine the biological properties of the final material required for use in vivo is described above and elsewhere (22).

**25 2. Materials and Methods**

Staphylokinase-neutralizing activity in plasma was determined as described above. Quantitation of antigen-specific IgG and IgM antibodies was performed using enzyme-linked immunosorbent assays in polystyrene  
30 microtiter plates essentially as described previously (22). In the IgG assays, dilution curves of affinospecific anti-SakSTAR IgG antibodies were included on each plate. These antibodies were isolated from plasma obtained from 3 patients, after thrombolytic therapy with  
35 wild-type SakSTAR, by chromatography on protein A-Sepharose and on insolubilized SakSTAR, and elution of bound antibodies with 0.1 mol/L glycine-HCl, pH 2.8. The purity of the IgG preparation was confirmed by sodium

007260 0640960

dodecylsulfate polyacrylamide gel electrophoresis. In the IgM assays, titers defined as the plasma dilution giving an absorbancy at 492 nm equivalent to that of a 1/640 dilution of pooled plasma were determined and compared with the titer of baseline samples before treatment (median value 1/410, interquartile range 1/120-1/700).

### 3. Thrombolytic efficacy

Wild-type SakSTAR or the variants SakSTAR(K74A) or SakSTAR(K74A,E75A,R77A) were administered intra-arterially at or in the proximal end of the occlusive thrombus as a bolus of 2 mg followed by an infusion of 1 mg/hr (reduced overnight in some patients to 0.5 mg/hr) in groups of 6 to 12 patients with angiographically documented occlusion of a peripheral artery or bypass graft of less than 120 days duration. Patients were studied after giving informed consent, and the protocol was approved by the Human Studies Committee of the University of Leuven. Inclusion and exclusion criteria, conjunctive antithrombotic treatment (including continuous intravenous heparin) and the study protocol were essentially as previously described (22).

Relevant baseline characteristics of the individual patients are shown in Table 2. The majority of PAO were at the femoropopliteal level. Two iliac stent and 8 graft occlusions were included. Eight patients presented with incapacitating claudication, 5 with chronic ischemic rest pain, 7 with subacute ischemia and 7 with acute ischemia. One patient (POE) who had 2 years previously been treated with SakSTAR was included in the SakSTAR(K74A) group. This patient was not included in the statistical analyses.

Table 2 also summarizes the individual treatment and outcome. Intra-arterial infusion, at a dose of 6.0 to 25 mg and a duration of 4.0 to 23 hrs, induced complete recanalization in 24 patients and partial recanalization in 3. Complementary endovascular procedures (mainly PTA) were performed in 17 patients and

001260 0640960



complementary reconstructive vascular surgery following thrombolysis in 3. No patient underwent major amputation. Early recurrence of thrombosis after the end of the angiographic procedure occurred in 4 patients. Bleeding complications were absent or limited to mild to moderate hematoma formation at the angiographic puncture sites except for 5 patients who required transfusion (data not shown). Intracranial or visceral hemorrhage was not observed. Circulating fibrinogen, plasminogen and  $\alpha_2$ -antiplasmin levels remained essentially unchanged during infusion of the SakSTAR moieties (data not shown), confirming absolute fibrin specificity of staphylokinase at the dosages used. Significant in vivo fibrin digestion occurred as evidenced by elevation of fibrin fragment D-dimer levels. Intra-arterial heparin therapy prolonged aPTT levels to a variable extent (data not shown).

#### 4. Antibody induction

Antibody-related SakSTAR-, SakSTAR(K74A)- and SakSTAR(K74A,E75A,R77A)-neutralizing activity and anti-SakSTAR, anti-SakSTAR(K74A) and anti-SakSTAR(K74A,E75A,R77A) IgG, were low at baseline and during the first week after the infusion (Figure 2). From the second week on, neutralizing activity levels increased to reach median values at 3 to 4 weeks of 20  $\mu$ g SakSTAR(K74A) and 2.4  $\mu$ g SakSTAR(K74A,E75A,R77A) neutralized per mL plasma in the patients treated with SakSTAR(K74A) and SakSTAR(K74A,E75A,R77A), respectively, which is significantly lower than the median value of 93  $\mu$ g wild-type SakSTAR neutralized per mL in the patients treated with SakSTAR ( $p=0.024$  for differences between the three groups by Kruskal-Wallis analysis and  $p=0.01$  and  $p=0.036$ , respectively, for variants vs wild-type by Mann-Whitney rank sum test). The levels of anti-SakSTAR(K74A) and of anti-SakSTAR(K74A,E75A,R77A) IgG increased to median values at 3 to 4 weeks of 270 and 82  $\mu$ g/mL plasma in patients treated with SakSTAR(K74A) and SakSTAR(K74A,E75A,R77A) respectively, which is

significantly lower than the median value of 1800  $\mu\text{g}$  anti-SakSTAR per mL plasma in the patients treated with SakSTAR (( $p= 0.024$  for differences between the three groups by Kruskal-Wallis analysis and  $p= 0.007$  and  $0.05$ , respectively, for variants versus wild-type by Mann-Whitney rank sum test).

The titers of anti-SakSTAR(K74A) and of anti-SakSTAR(K74A,E75A,R77A) IgM increased from median baseline values of 1/460 and 1/410 to median values at 1 week of 1/510 and 1/450 in patients treated with SakSTAR(K74A) and SakSTAR(K74A,E75A,R77A), respectively, which was not significantly different from the median values of 1/320 at baseline and 1/640 at week 1 in patients treated with SakSTAR. Corresponding values at 2 weeks were 1/590 and 1/550 in patients given SakSTAR(K74A) and SakSTAR(K74A,E75A,R77A), not significantly different from 1/930 with SakSTAR (data not shown). The antibodies induced by treatment with SakSTAR were completely absorbed by SakSTAR but incompletely by SakSTAR(K74A) and by SakSTAR(K74A,E75A,R77A) confirming the immunogenicity of the K74,E75,R77 epitope and the dominant role of K74 in the binding of antibodies directed against this epitope. The antibodies induced by treatment with SakSTAR(K74A) or SakSTAR(K74A,E75A,R77A) were completely absorbed by SakSTAR, by SakSTAR(K74A) and by SakSTAR(K74A,E75A,R77A), indicating that immunization was not due to neoepitopes generated by substitution of Lys74 with Ala, but to epitopes different from the K74,E75,R77 epitope.

Thus, this example illustrates that staphylokinase variants with reduced antibody induction but intact thrombolytic potency can be generated. The present experience in 26 patients treated with SakSTAR ( $n= 9$ ), SakSTAR(K74A) ( $n= 11$ ) and SakSTAR(K74A,E75A,R77A) ( $n= 6$ ) combined with previous experience in 14 patients with SakSTAR ( $n= 7$ ) and SakSTAR(K35A, E38A,K74A,E75A,R77A) ( $n= 7$ ) (31) and in 24 patients with SakSTAR (32), and with subsequent non-randomized

experience in patients with SakSTAR (n= 30) with SakSTAR(K74A) (n= 12) and with SakSTAR(K74A,E75A,R77A) (n= 7) (data not shown), allows an initial estimation of the prevalence of immunization by intra-arterial treatment with SakSTAR or variants with an altered K74,E75,R77 epitope [SakSTAR(K74A), SakSTAR(K74A,E75A,R77A) and SakSTAR(K35A,E38A,K74A,E75A, R77A)]. Neutralizing activity data after 2 to 4 weeks, available in 70 patients with peripheral arterial occlusion given intra-arterial SakSTAR, revealed that 56 patients (80 percent) had levels > 5 µg compound neutralized per mL plasma. Of the patients given SakSTAR(K74A), SakSTAR(K74A,E75A, R77A) or SakSTAR(K74A,E75A,K74A,E75A, R77A), 27 of the 43 (63 percent) had neutralizing activity levels of > 5 µg compound per mL plasma. This difference is statistically significant (p= 0.05 by Fisher's exact test) indicating that the K74,E75,R77 epitope is a major determinant of antibody induction.

#### 20 **EXAMPLE 4**

Construction, epitope mapping with murine monoclonal antibodies and absorption with pooled plasma of immunized patients, of alanine-substitution mutants of staphylokinase

##### 25 1. Introduction

Site-directed mutagenesis was applied to residues other than "charged amino acids" in order to identify i) additional residues belonging to epitopes I and III identified with the panel of murine Mabs and ii) amino acids determining absorption to antiserum from immunized patients. Since functional epitopes generally comprise more than one amino acid residue critical for antibody binding, identification of additional residues in these epitopes could lead to the construction of new combination derivatives displaying a lower antigenic profile, while keeping the specific activity and the temperature stability of wild-type staphylokinase. In this example, the construction and characterization of

001260 06470960

SakSTAR variants in which one or at most two amino acids (adjacent or in close vicinity) were substituted with alanine is described. The mutants described under this example are listed in Table 3. These variants were

- 5 expressed in E. coli, purified and characterized in terms of specific activity, reactivity with the panel of murine monoclonal antibodies, and absorption of antibodies from plasma of patients treated with wild-type SakSTAR.

10 2. Reagents and Methods

The source of all reagents used in the present study has previously been reported (22), or is specified below. The template vector for mutagenesis, pMEX602sakB (i.e. pMEX.SakSTAR), has been described elsewhere (23).

- 15 Restriction and modification enzymes were purchased from New England Biolabs (Leusden, The Netherlands), Boehringer Mannheim (Mannheim, Germany) or Pharmacia (Uppsala, Sweden). The enzymatic reactions were performed according to the supplier recommendation. The mutagenic  
20 oligonucleotides and primers were obtained from Eurogentec (Seraing, Belgium). Plasmid DNA was isolated using a purification kit from Qiagen (Hilden, Germany) or the BIO 101 RPM kit (Vista, CA), as recommended. Transformation-competent E. coli cells were prepared by  
25 the well-known calcium phosphate procedure. Nucleotide sequence determination was performed on double strand plasmid DNA with the dideoxy chain termination method, using the T7 sequencing kit (Pharmacia, Uppsala, Sweden). Polymerase chain reactions (PCR) were performed using Taq  
30 polymerase from Boehringer Mannheim (Mannheim, Germany) or Vent polymerase (New England Biolabs, Leusden, The Netherlands). The recombinant DNA methods required to construct the variants described in this example are well established (22, 27).

35

3. Construction of expression plasmids

The variants SakSTAR(Y17A,F18A), Sak-STAR(F104A), SakSTAR(F111A), SakSTAR(Y9A), SakSTAR(Y91A),

007250 0640960

The variant SakSTAR(F125A) was constructed by performing two consecutive PCR reactions (30 cycles: : 30 sec at 94°C, 30 sec at 50°C, 30 sec at 72°C). In the first reaction, a fragment of pMEX.SakSTAR was amplified  
35 with the primers 818A and a mutagenic primer. This amplified fragment was then used as template in a second PCR reaction with a mutagenic primer in order to further elongate the fragment downstream of the StyI site present

in the sakSTAR gene (corresponding to amino acids 130-131 of SakSTAR). The resulting product was digested with EcoRI and StyI, and ligated into the corresponding sites of pMEXSakSTAR.

5           The plasmids encoding all the other variants listed in Table 3 were constructed by direct PCR or by the spliced overlap extension polymerase chain reaction (SOE-PCR) (24) using pMEX.SakSTAR or available plasmids encoding SakSTAR variants as template. Two fragments were  
10 amplified by PCR (30 cycles: 1 sec at 94°C, 1 sec at 50°C, 10 sec at 72°C), the first one starting from the 5' end (primer 818A ) of the staphylokinase gene to the region to be mutagenized (forward primer), the second one from this same region (backward primer) to the 3' end of  
15 the gene with primer 818D (5' CAAACAGCCAAGCTTCATTCATTCAGC). The forward and backward primers shared an overlap of around 24 bp (primers not shown). The two purified fragments were then assembled together in a second PCR reaction with the  
20 external primers 818A and 818D (30 cycles: 1 sec at 94°C, 1 sec at 50°C, 10 sec at 72°C). The amplified product from this final reaction was purified, digested with EcoRI and HindIII and ligated into the corresponding site of pMEX.SakSTAR. For each construction, the sequence of  
25 the variant was confirmed by sequencing the entire SakSTAR coding region.

#### 4. Expression and purification of SakSTAR variants

30           The SakSTAR variants were expressed and purified, as described below, from transformed E. coli grown in terrific broth (TB) medium (28). A 2 to 4 mL aliquot of an overnight saturated culture in LB medium was used to inoculate a 1 to 2 L culture in terrific broth supplemented with 100 µg/mL ampicillin. The culture  
35 was incubated with vigorous aeration and at 30°C. After about 16 hours incubation, IPTG (200 µmol/L) was added to the culture to induce expression from the tac promoter. After 3 hours induction, the cells were pelleted by

007260 0640960

centrifugation at 4,000 rpm for 20 min, resuspended in 1/10 volume of 0.01 mol/L phosphate buffer pH 6-6.5 and disrupted by sonication at 0°C. The suspension was centrifuged for 20 min at 20,000 rpm and the supernatant was stored at 4°C or at -20°C until purification. The material was purified essentially as described above (Example 2): cleared cell lysates containing the SakSTAR variants were subjected to chromatography on a 1.6 x 5 cm column of SP-Sephadex, followed by chromatography on a 1.6 x 8 cm column of phenyl-Sepharose. The SakSTAR containing fractions, localized by SDS-gel electrophoresis, were pooled for further analysis.

#### 5. Physicochemical and biochemical analysis

Protein concentrations were determined according to Bradford (29). SDS-PAGE was performed with the Phast System<sup>TM</sup> (Pharmacia, Uppsala, Sweden) using 10-15% gradient gels and Coomassie Brilliant blue staining, and the specific activities of SakSTAR solutions were determined with a chromogenic substrate assay carried out in microtiter plates (as described in example 2). The specific activity of the different SakSTAR variants are summarized in Table 3.

#### 6. Reactivity of SakSTAR variants with a panel of murine monoclonal antibodies

The methodology used to determine the reactivity of the SakSTAR variants with a panel of murine monoclonal antibodies was described in example 1 above. The results are summarized in Table 3 (the layout of this Table corresponds to the layout of Table 1, as described in example 1). Apparent association constants at least 10-fold lower than those of wild-type staphylokinase were considered as significant and are indicated in bold type in the table.

In order to obtain a comprehensive inventory of the properties of Ala-substitution variants of the SakSTAR molecule, 67 plasmids encoding variants with

substitution of a single or two adjacent amino acids with Ala were constructed, expressed and purified. Together with the 35 charged residue to Ala-substitution variants previously described (22, and example 2), this analysis covers all residues in SakSTAR except Gly, Ala and Pro, as illustrated in Figure 3. Eight of the variants could not be obtained in purified form, primarily as a result of low expression levels, 11 variants were inactive, 56 had a reduced specific activity, and 27 had a maintained or increased specific activity ( $\geq 100$  kHU/mg). The yields of purified material from cultures of expressed plasmids were 16 mg/L (median, 10 to 90 percentile range 4 to 41 mg/L). SDS polyacrylamide gel electrophoresis consistently showed one main band with  $M_r \approx 16,000$ , usually representing 95% of total protein (not shown).

Substitution of K35, N95, S103 or K135 with Ala yielded variants with specific activities of  $\geq 200$  kU/mg. Substitution of W66, Y73 or E75 with Ala reduced the reactivity of the variants with  $\geq 3$  antibodies of epitope cluster I, of H43 or V45 with Ala that with 3 antibodies from epitope cluster II and of V32, K35, D82 and K130 with Ala that with  $\geq 3$  antibodies of epitope cluster III.

7. Absorption of antibodies, elicited in patients by treatment with SakSTAR

For the present example, the three plasma pools, as described in example 2 were used. The methodology used to evaluate the absorption with wild-type staphylokinase and with SakSTAR variants, of antibodies elicited in patients treated with SakSTAR, is described in detail in example 2. The results are summarized in Table 3. Whereas wild-type SakSTAR and most of the variants analyzed in this example absorbed more than 95% of the binding antibodies from pooled plasma of the 10 patients, incomplete absorption ( $< 60\%$ ) was observed with SakSTAR(Y73A), and with SakSTAR(K74A). The predominant role of Lys74 for antibody recognition has



been demonstrated previously (see example 2). The present results indicate that Tyr73 participates to the same major epitope as Lys74, or, alternatively, that substitution at Tyr73 may indirectly induce a structural  
5 modification of the "K74-epitope". Absorption with pooled plasma from 3 patients from which >95% of the antibodies were absorbed with SakSTAR(K35A,E38A,K74A,E75A,R77A) (Subpool C, see example 2) was nearly complete with most variants tested.

10

**EXAMPLE 5**

Construction, epitope mapping with murine monoclonal antibodies and absorption with pooled plasma of immunized patients, of staphylokinase variants with substitution of  
15 S34, G36 and/or H43

The natural variant Sak42D differs from SakSTAR in three amino acids and corresponds to SakSTAR(S34G, G36R,H43R). Sak42D is characterized by reduced reactivity with some murine antibodies of epitope clusters II and  
20 III and a slightly reduced absorption of antibodies from plasma of patients treated with SakSTAR (Table 4). Mutagenesis of these residues in SakSTAR revealed that the reduced reactivity with epitope cluster III and with immunized patient plasma could be ascribed to the G36R  
25 substitution, the H43R substitution mediated the reduced reactivity with epitope cluster II but had no effect on the reactivity with immunized patient plasma, whereas the S34A substitution had no effect. The G36R substitution could be combined with the K74R but not with the K74A  
30 substitution, without significant reduction of the specific activity (Table 4).

001260"06410960

**EXAMPLE 6**

Construction, epitope mapping with murine monoclonal antibodies and absorption with pooled plasma of immunized patients, of staphylokinase variants with substitution of

5 K35, E65, Y73, K74, E80+D82, N95, K130, V132 and/or K135

Based on the results of the alanine-substitution analysis in example 4, K35, N95 and K135 were selected for further analysis because SakSTAR(K35A), SakSTAR(N95A) and SakSTAR(K135A) had a two-fold increased  
10 specific activity, Y73 and K74 because SakSTAR(Y73A) and SakSTAR(K74A) had a markedly reduced reactivity with antibodies from epitope cluster I and diminished absorption of antibodies from plasma of patients immunized by treatment with SakSTAR, and K35, E80+D82,  
15 K130 and V132 because SakSTAR(K35A), SakSTAR(E80A,D82A), SakSTAR(K130A) and SakSTAR(V132A) had a reduced reactivity with antibodies from epitope cluster III.

In an effort to maximize the activity/antigenicity ratio, these amino acids were substituted  
20 with other amino acids than Ala. As summarized in Table 5, substitution of K35 with A, E or Q revealed that SakSTAR(K35A) had the most interesting properties, substitution of Y73 with F, H, L, S or W did not rescue the marked reduction in specific activity, and K74  
25 confirmed its key role in binding of antibodies from immunized patient plasma, the best specific activity/antigenicity ratios being obtained with SakSTAR(K74Q) and SakSTAR(K74R). SakSTAR(E80A,D82A) was preferred over the single residue variants SakSTAR(E80A) or SakSTAR(D82A)  
30 because of its somewhat lower reactivity with immunized patient plasma. SakSTAR(N95A) could not be further improved by substitution of N95 with E, G, K or R and it was unable to confer its increased specific activity to variants containing K74A or K135R. Finally SakSTAR(K130A)  
35 was outperformed in terms of specific activity by SakSTAR(K130T) and SakSTAR(V132A) by SakSTAR(V132R).

007250 06470960

**EXAMPLE 7**

Construction, epitope mapping with murine monoclonal antibodies and absorption with pooled plasma of immunized patients of combination variants of SakSTAR(K130T,K135R)  
5 and SakSTAR(E80A,D82A,K130T,K135R) with  
K35A,G36R,E65X,K74X and selected other amino acids

In the present and the following examples an additional plasma pool was made from 40 patients obtained several weeks after treatment with SakSTAR (Pool 40). The  
10 original pool from 10 patients is further identified as Pool 10. The absorption of staphylokinase-specific antibodies was quantified as described above and elsewhere (22).

The SakSTAR(K130T,K135R) variant was taken as a  
15 template for additive mutagenesis because of its high specific activity with a moderate reduction of binding to antibodies of epitope cluster III and absorption of antibodies from immunized patient plasma (Table 6). Addition of G36R, K74R, or K74Q or both to the template  
20 did not markedly reduce the specific activity, reduced the reactivity with monoclonal antibodies against epitope cluster III (G36R substitution) and decreased the absorption of antibodies from immunized patient plasma (K74R or K74Q substitution). Combination of E65A or E65Q  
25 with K74Q in the SakSTAR(K130T,K135R) template reduced the absorption of antibodies from Pool 10 and Pool 40 to around 50 and 60 percent respectively, without markedly reducing the specific activity. Addition substitution of selected amino acids in the SakSTAR(E65Q,K74Q,K130T,  
30 K135R) template did not further reduce the antibody absorption from Pool 10 or Pool 40. Surprisingly, the substitution of K136 with A and the addition of K in position 137 resulted in a marked increase in specific activity, as measured in the chromogenic substrate assay.  
35 Combination of the SakSTAR(E80A,D82A) and SakSTAR(K130T,K135R) templates, did not affect the specific activity and had a reduced reactivity with epitope cluster III antibodies (Table 7). Therefore the Sak-

09601490.092400

STAR(E80A,D82A,K130T,K135R) template was selected for further mutagenesis. Addition of K74R and even more of K74Q drastically reduced the reactivity with immunized patient plasma. Finally, addition of E65D or of K35A or  
 5 E65S to the SakSTAR(K74R,E80A,D82A,K130T,K135R) or Sak-STAR(K74Q,E80A,D82A,K130T, K135R) templates yielded variants with intact specific activity which only bound  
 ≤45 of the antibodies of pooled immunized patient plasma and less than 15 percent of the subpool reacting for more  
 10 than 50 percent with the K74,E75,R77 epitope.

#### EXAMPLE 8

Characterization of selected variants of staphylokinase with intact specific activity and less than 50%

15 adsorption of pooled SakSTAR specific human antibodies elicited in patients by treatment with wild-type SakSTAR

##### 1. Introduction

Twenty three of the variants constructed and characterized in the above examples combined the  
 20 properties of a residual specific activity of ≥100 KHU/mg and ≤50 percent absorption with the pool of antisera obtained from 10 patients treated with wild-type SakSTAR. The results are summarized in Table 8. Results obtained with Subpool B and Subpool C and with the pool of 40  
 25 patients treated with wild-type SakSTAR are included. SakSTAR(K74Q,E80A,D82A,K130T, K135R), SakSTAR(E65D,K74R, E80A,D82A,K130T,K135R), SakSTAR(K35A,E65D,K74Q, E80A, D82A,K130T,K135R) and SakSTAR(E65Q,K74Q,N95A,E118A, K130A,K135R,K136A,V137K) were selected for further  
 30 characterization.

##### 2. Fibrinolytic properties of SakSTAR variants in human plasma in vitro

The fibrinolytic and fibrinogenolytic  
 35 properties of the SakSTAR variants were determined as previously described. Dose- and time-dependent lysis of <sup>125</sup>I-fibrin labeled human plasma clots submerged in human plasma was obtained with the selected variants (Table 9).

007260" 05470960

Spontaneous clot lysis during the experimental period was  $\leq 5\%$  (not shown). Equi-effective concentrations of test compound (causing 50% clot lysis in 2 hrs;  $C_{50}$ ), determined graphically from plots of clot lysis at 2 hrs versus the concentration of plasminogen activator (not shown), ranged from  $0.11 \pm 0.01$  to  $0.24 \pm 0.04$  g/mL at which the residual fibrinogen levels ranges between  $92 \pm 30$  and  $97 \pm 30$  percent of baseline (Table 9). The concentrations of compound causing 50% fibrinogen degradation in 2 hrs in human plasma in the absence of fibrin were determined graphically from dose-response curves (not shown). These values (mean  $\pm$  SD of 3 independent experiments) ranged from  $14 \pm 3.2$  to  $29 \pm 3.1$   $\mu$ g/mL (Table 9). Surprisingly the very high specific activity of SakSTAR(E65Q,K74Q,N95A,E118A,K130A,K135R,K136A,V137K) in the chromogenic assay was not associated with an increased thrombolytic potency in a plasma milieu.

3. Temperature stability of selected SakSTAR variants

The temperature stability of preparations of SakSTAR(K74Q,E80A,D82A,K130T,K135R), SakSTAR(E65D,K74R,E80A,D82A,K130T,K135R) and SakSTAR(K35A,E65D,K74Q,E80A,D82A,K130T,K135R), dissolved to a concentration of 1.0 mg/mL in 0.15 mol/L NaCl, 0.01 mol/L phosphate buffer, pH 7.5 at various temperatures is illustrated in Fig. 4. At temperatures up to 37°C, all compounds remained fully active for up at least three days. At 56°C and 70°C the three variants were however less stable than wild-type SakSTAR.

4. Pharmacokinetic properties of SakSTAR variants following bolus injection in hamsters

The pharmacokinetic parameters of the disposition of SakSTAR variants from blood were evaluated in groups of 4 hamsters following intravenous bolus injection of 100  $\mu$ g/kg SakSTAR variant. SakSTAR-related antigen was assayed using the ELISA described elsewhere.

The ELISA was calibrated against each of the SakSTAR variants to be quantitated. Pharmacokinetic parameters included: initial half-life (in min),  $t_{1/2\alpha} = \ln 2/\alpha$ ; terminal half-life (in min),  $t_{1/2\beta} = \ln 2/\beta$ ; volume of the central (plasma) compartment (in mL),  $V_c = \text{dose}/(A+B)$ ; area under the curve (in  $\mu\text{g}\cdot\text{min}\cdot\text{mL}^{-1}$ ),  $\text{AUC} = A/\alpha + B/\beta$ ; and plasma clearance (in  $\text{mL}\cdot\text{min}^{-1}$ ),  $\text{Clp} = \text{dose}/\text{AUC}$  (33).

The disposition rate of staphylokinase-related antigen from blood following bolus injection of 100  $\mu\text{g}/\text{kg}$  of the selected SakSTAR variants in groups of 4 hamsters could adequately be described by a sum of two exponential terms by graphical curve peeling (results not shown), from which the pharmacokinetic parameters summarized in Table 10 were derived. The pharmacokinetic parameters of the mutants were not markedly different from those of wild type SakSTAR. Initial plasma half-lives ( $t_{1/2}(\alpha)$ ) ranged between 2.0 and 3.2 min and plasma clearances (Clp) between 1.6 and 4.1 mL/min.

## 20 EXAMPLE 9

Comparative thrombolytic efficacy and immunogenicity of SakSTAR(K74Q,E80A,D82A, K130T,K135R) and SakSTAR(E65D,K74R,E80A,D82A,K130T,K135R) versus SakSTAR in patients with peripheral arterial occlusion

### 25 1. Purification for use in vivo

Eighteen liter cultures (in 2 L batches) of the variants SakSTAR(K74Q,E80A,D82A,K130T, K135R) and SakSTAR(E65D,K74R,E80A,D82A,K130T,K135R) were grown for 20 hours in terrific broth medium (28), supplemented with 100  $\mu\text{g}/\text{mL}$  ampicillin and induced with IPTG during the last 3 hours. The cells were pelleted, resuspended in 1/10 volume of 0.01 mol/L phosphate buffer, pH 6.0, disrupted by sonication and cleared by centrifugation. The compounds were purified by chromatography on a 10 x 7 cm column of SP-Sepharose, equilibrated with 0.01 mol/L phosphate buffer, pH 6.0 and eluted with a 1 mol/L NaCl gradient (3 column volumes). The fractions containing SakSTAR variant were pooled, solid NaCl was added to a

concentration of 2.5 mol/L and the material was chromatographed on a 10 x 20 cm column of phenyl-Sepharose followed by stepwise elution with 0.01 mol/L phosphate buffer, pH 6.0. The materials were  
5 desalted on a 10 x 45 cm column of Sephadex G25, concentrated by application on a 5 x 10 cm column of SP-Sepharose with stepwise elution with 1.0 mol/L NaCl and finally gel filtered on a 6 x 60 cm column of Superdex 75 equilibrated with 0.15 m NaCl, 0.01 mol/L  
10 phosphate buffer, pH 7.5 to further reduce their endotoxin content. The SakSTAR variant containing fractions were pooled, the protein concentration was adjusted to 1 mg/mL and the material sterilized by filtration through a 0.22 m Millipore filter. The  
15 methodology used to determine specific activity, endotoxin contamination, bacterial sterility and toxicity in mice is described above and elsewhere (22). The purity of the preparation was evaluated by SDS gel electrophoresis on 10% gels to which 40 g of compound was  
20 applied.

Out of culture volumes of 18 liters of SakSTAR variant, 840 mg of SakSTAR(K74Q,E80A, D82A,K130T,K135R) with a specific activity of 140 kHU/mg and 800 mg Sak-STAR(E65D, K74R,E80A,D82A,K130T,K135R) with a specific  
25 activity of 150 were purified. The endotoxin content was <0.1 and 0.26 IU/mg. Gel filtration on HPLC revealed a single main symmetrical peak in the chromatographic range of the column, representing >98% of the eluted material (total area under the curve) (not shown). SDS gel  
30 electrophoresis of 40 g samples revealed single main components (not shown). Preparations sterilized by filtration proved to be sterile on 3 day testing as described elsewhere (22). Intravenous bolus injection of SakSTAR variants in groups of 5 mice (3 mg/kg body  
35 weight), did not provoke any acute reaction, nor reduced weight gain within 8 days, in comparison with mice given an equal amount of saline (not shown).

001250-05470960

## 2. Thrombolytic efficacy

Wild-type SakSTAR or the variants SakSTAR(K74Q, E80A,D82A,K130T,K135R) or SakSTAR(E65D,K74R,E80A,D82A, K130T,K135R) were administered intra-arterially at or in the proximal end of the occlusive thrombus as a bolus of 2 mg followed by an infusion of 1 mg/hr (reduced overnight in some patients to 0.5 mg/hr) in groups of 15, 6 and 6 patients respectively with angiographically documented occlusion of a peripheral artery or bypass graft of less than 30 days duration. Patients were studied after giving informed consent, and the protocol was approved by the Human Studies Committee of the University of Leuven. Inclusion and exclusion criteria, conjunctive antithrombotic treatment (including continuous intravenous heparin) and the study protocol were essentially as previously described (22).

Relevant baseline characteristics of the individual patients and results of treatment and outcome are shown in Table 11. Intra-arterial infusion, at a dose of 3.5 to 27 mg and a duration of 2 to 44 hrs, induced complete recanalization in 22 patients and partial recanalization in 5. Complementary endovascular procedures (mainly PTA) were performed in 13 patients and complementary reconstructive vascular surgery following thrombolysis in 5. One patient underwent major amputation. Bleeding complications were usually absent or limited to mild to moderate hematoma formation at the angiographic puncture sites (data not shown). One patient, given wild-type SakSTAR suffered a non-fatal intracranial bleeding, one (BUE) a retroperitoneal hematoma and two (MAN and STRO) a gastro-intestinal bleeding.

Circulating fibrinogen, plasminogen and  $\alpha_2$ -antiplasmin levels remained unchanged during infusion of the SakSTAR moieties (data not shown), reflecting absolute fibrin specificity of these agents at the dosages used (data not shown). Significant in vivo fibrin digestion occurred as evidenced by elevation of fibrin

007260-06470960



fragment D-dimer levels. Intra-arterial heparin therapy prolonged aPTT levels to a variable extent (data not shown).

5 3. Antibody induction

Staphylokinase-neutralizing activity in plasma and antigen-specific IgG antibodies were quantitated essentially as described above and elsewhere (22). Antibody-related SakSTAR-, SakSTAR(K74Q,E80A,D82A, K130T,K135R)- and SakSTAR(E65D,K74R,E80A,D82A, K130T,K135R)-neutralizing activity and anti-SakSTAR, anti-SakSTAR(K74Q,E80A,D82A,K130T,K135R) and anti-SakSTAR(E65D,K74R,E80A, D82A,K130T,K135R) IgG, were low at baseline and during the first week after the infusion (Figure 5). From the second week on, neutralizing activity levels increased to reach median values at 3 to 4 weeks of 9  $\mu$ g SakSTAR(K74Q,E80A,D82A, K130T,K135R) and 0.5  $\mu$ g SakSTAR(E65D,K74R,E80A,D82A, K130T,K135R) neutralized per mL plasma in the patients treated with the corresponding moieties, respectively, as compared to median value of 24  $\mu$ g wild-type SakSTAR neutralized per mL in the 15 patients treated with SakSTAR. The levels of anti-SakSTAR(K74Q,E80A,D82A, K130T,K135R) and of anti-SakSTAR(E65D,K74R,E80A, D82A,K130T,K135R) IgG increased to median values at 3 to 4 weeks of 420 and 30  $\mu$ g/mL plasma in patients treated with the corresponding moieties, respectively, as compared to a median value of 590  $\mu$ g anti-SakSTAR per mL plasma in the patients treated with SakSTAR (Figure 5). The prevalence of immunization, defined as neutralizing activities in plasma after 2 to 4 weeks exceeding 5 g/ml was 3 of 6 patients (50 percent) with SakSTAR(K74Q,E80A, D82A,K130T,K135R), 1 of 6 patients (17 percent) with SakSTAR(E65D,K74R,E80A,D82A, K130T,K135R), as compared to 56 of 70 patients (80 percent) with SakSTAR. This difference is statistically highly significant ( $p= 0.01$  by 2 x 3 Chi square analysis).

The antibodies induced by treatment with SakSTAR were completely absorbed by SakSTAR but incompletely by SakSTAR(K74Q,E80A,D82A,K130T,K135R) and by SakSTAR(E65D,K74R, E80A,D82A,K130T,K135R) (Table 12).

- 5 Antibodies induced by treatment with Sak-STAR(K74Q,E80A,D82A,K130T,K135R), detectable in 4 of the 6 patients, were completely ( $\geq 90$  percent) absorbed by SakSTAR, by SakSTAR(K74Q,E80A,D82A,K130T, K135R) and by SakSTAR(E65D,K74R,E80A,D82A,K130T,K135R), indicating that
- 10 immunization was not due to neoepitopes generated by substitution of wild-type amino acids. Antibodies induced by treatment with SakSTAR(E65D,K74R,E80A,D82A, K130T,K135R) detectable in one patient (URB) were completely absorbed with SakSTAR(K74Q,E80A,D82A,
- 15 K130T,K135R) and with SakSTAR(E65D,K74Q,E80A,D82A, K130T,K135R) but incompletely (85%) with wild-type SakSTAR, suggesting that a small fraction of the induced antibodies might be directed against a neoepitope in the variant used for infusion.

20

#### EXAMPLE 10

Construction and absorption with pooled plasma of immunized patients of combination variants of SakSTAR(E65Q,K74Q,K130T,K135R) and other selected amino

25 acids

##### 1. Introduction

- In a final round of additive substitution mutagenesis, the SakSTAR(E65Q,K74Q,K130T, K135R) variant was taken as a template because it displayed a high
- 30 specific activity with a significant reduction of absorption (to 65 percent) of antibodies from pooled immunized patient plasma (Pool 40). The intermediate variants which were relevant for the composition of the finally selected variants are summarized in Table 13.
- 35 Addition of K35A, D82A and S84A, of T90A,E99D and T101S or of E108A and K109A reduced the antibody absorption to around 50 percent, whereas the combined addition of D82A,S84A and E108A, K109A reduced it to 41 percent.

007260"064T0960

Substitution of K136A combined with the addition of a Lys at the COOH terminus (-137K) increased the specific activity in a purified system but not in a plasma milieu nor in a hamster pulmonary embolism model (not shown), and further reduced the absorption of antibodies from pooled patient plasma to 30 percent. Finally, addition of the K35A, and T90A, E99D, T101S substitutions to this template yielded a mutant with intact thrombolytic potency which only bound 24 percent of the antibodies of pooled immunized patient plasma.

Based on this analysis, SakSTAR(E65Q, K74Q, D82A, S84A, E108A, K109A, K130T, K135R, K136A, V137K), (SY118), and SakSTAR(K35A, E65Q, K74Q, D82A, S84A, T90A, E99D, T101S, E108A, K109A, K130T, K135R, K136A, V137K), (SY141), were selected for further characterization. In addition, SakSTAR(K35A, E65Q, K74R, D82A, S84A, T90A, E99D, T101S, E108A, K109A, K130T, K135R, K136A, V137K), (SY145) with a Lys in position 74, was constructed and evaluated.

## 2. Pharmacokinetic properties of SakSTAR variants following bolus injection in hamsters

The disposition rate of staphylokinase-related antigen from blood following bolus injection of 100 µg/kg of the selected SakSTAR variants in groups of 4 hamsters could adequately be described by a sum of two exponential terms by graphical curve peeling (results not shown). The pharmacokinetic parameters of the mutants were derived from these plasma disappearance curves not markedly different from those of wild type SakSTAR (results very similar to those of table 10, data not shown).

### EXAMPLE 11

#### Characterization of selected variants derived from SakSTAR(E65Q, K74Q, K130T, K135R)

## 1. Fibrinolytic properties of selected SakSTAR variants towards human plasma in vitro

Dose- and time-dependent lysis of <sup>125</sup>I-fibrin labeled human plasma clots submerged in human plasma was

092400-092400

5 determined graphically from plots of clot lysis at 2 hrs  
versus the concentration of plasminogen activator (not  
shown), ranged from  $0.15 \pm 0.02$  to  $0.19 \pm 0.01$   $\mu\text{g/ml}$  at  
which no significant fibrinogen degradation occurred. The  
concentrations of compound causing 50% fibrinogen  
10 degradation in 2 hrs in human plasma in the absence of  
fibrin were determined graphically from dose-response  
curves (not shown). These values (mean  $\pm$  SD of 3  
independent experiments) ranged from  $7.0 \pm 0.6$  to  $24 \pm$   
 $3.6$   $\mu\text{g/ml}$  (Table 14).

## 15

30

**EXAMPLE 12**

Comparative thrombolytic efficacy and immunogenicity of SakSTAR(E65Q,K74Q, D82A,S84A,E108A,K109A,K130T,K135R, K136A,V137K), (SY118), SakSTAR(K35A, E65Q,K74Q,D82A,S84A, T90A,E99D,T101S,E108A,K109A,K130T,K135R,K136A, VV137K), (SY141), and SakSTAR(K35A,E65Q,K74R,D82A,S84A,T90A, E99D,T101S, E108A,K109A,K130T,K135R,K136A,V137K), (SY145), in patients with peripheral arterial occlusion

1. Large scale purification and conditioning of SakSTAR variants for use in vivo

Material was purified to homogeneity out of culture volumes of 18 liters. The endotoxin content was below 2 IU/mg. Gel filtration on HPLC revealed a single main symmetrical peak in the chromatographic range of the column, representing >98% of the eluted material (total area under the curve) (not shown). SDS gel electrophoresis of 30 µg samples revealed single main components. Preparations sterilized by filtration proved to be sterile on 3 day testing. Intravenous bolus injection of SakSTAR variants in groups of 5 mice (3 mg/kg body weight), did not provoke any acute reaction, nor reduced weight gain within 8 days, in comparison with mice given an equal amount of saline (not shown).

Groups of 6 patients with angiographically documented peripheral arterial occlusion (PAO) were studied. Relevant baseline characteristics of the individual patients are shown in Table 15. Table 16 summarizes the individual treatment and outcome. Intra-arterial infusion, at a dose of 6 to 24 mg and a duration of 4 to 29 hrs, induced complete recanalization in most patients. Circulating fibrinogen, plasminogen and  $\alpha_2$ -antiplasmin levels remained essentially unchanged during infusion of the SakSTAR variants (data not shown), reflecting absolute fibrin specificity of these agents at the dosages used. Antibody-related SakSTAR(E65Q,K74Q, D82A,S84A,E108A,K109A,K130T,K135R,K136A,V137K)-, Sak-STAR(K35A,E65Q,K74Q,D82A,S84A,T90A,E99D,T101S,E108A,

002250" 06410950

K109A, K130T, K135R, K136A, V137K) - and SakSTAR(K35A, E65Q, K74R, D82A, S84A, T90A, E99D, T101S, E108A, K109A, K130T, K135R, K136A, V137K) - neutralizing activity, were low at baseline and during the first week after the infusion (Table 17).

- 5 From the second week on neutralizing activity levels increased to reach median values at 3 to 4 weeks of 19  $\mu$ g SakSTAR(E65Q, K74Q, D82A, S84A, E108A, K109A, K130T, K135R, K136A, V137K), (SY118), 0.7  $\mu$ g SakSTAR(K35A, E65Q, K74Q, D82A, S84A, T90A, E99D, T101S, E108A, K109A, K130T, K135R, K136A, V137K), (SY141), and 4.3  $\mu$ g SakSTAR(K35A, E65Q, K74R, D82A, S84A, T90A, E99D, T101S, E108A, K109A, K130T, K135R, K136A, V137K), (SY145), neutralized per ml plasma in the patients treated with the respective compounds, which for SY141 and SY145, but not for SY118 is lower than the
- 15 median value of 12  $\mu$ g wild type SakSTAR neutralized per ml in 69 patients treated with wild type SakSTAR.

- Overt immunization (neutralizing activity at 3 to 4 weeks of 5 g compound per ml plasma) was observed in 56 of 70 patients treated with SakSTAR, in 5 of the 6
- 20 patients exposed to SakSTAR(E65Q, K74Q, D82A, S84A, E108A, K109A, K130T, K135R, K136A, V137K), (SY118), only in 2 of the 6 patients given SakSTAR(K35A, E65Q, K74Q, D82A, S84A, T90A, E99D, T101S, E108A, K109A, K130T, K135R, K136A, V137K), (SY141), and in 1 of the 3 patients given SakSTAR(K35A, E65Q, K74R, D82A, S84A, T90A, E99D, T101S, E108A, K109A, K130T, K135R, K136A, V137K), (SY145).
- 25

- The results with respect to immunogenicity of the main variants studied in patients are summarized in Table 18. Clearly, variants SakSTAR(E65D, K74R, E80A, D82A, K130T, K135R) and SakSTAR(K35A, E65Q, K74Q, D82A, S84A, T90A, E99D, T101S, E108A, K109A, K130T, K135R, K136A, V137K) have a significantly reduced immunogenicity when compared to the wild type protein.
- 30

**EXAMPLE 13****Construction, purification and characterization of cysteine-substitution mutants of staphylokinase****1. Introduction**

5           Site-directed mutagenesis was applied to substitute exposed amino acids with single cysteine residues in order to construct i) homodimeric forms of staphylokinase, upon formation of an intermolecular disulfide bridge, and ii) polyethylene glycol-conjugated  
10 molecules (PEG-derivatives). The aim of this example was twofold: first, the clearance can be reduced by increasing the size of the injected molecule (via dimerization or conjugation with large molecule such as PEG) and second, PEG-derivatives have also been shown to  
15 induce a reduced immunoreactivity in animal models (for review, see ref. 34). In both cases, a prolonged half-life in vivo could help to reduce the pharmacological dose of staphylokinase in patients. This reduction could be accompanied with a reduced immunogenic  
20 reaction against the thrombolytic agent, thus enhancing its pharmacological activity as a thrombolytic agent.

          In this example, the construction and characterization of two SakSTAR variants in which one single amino acid was substituted with cysteine is  
25 described. The mutants described under this example are listed in Table 19. These variants were expressed in E. coli, purified and characterized in terms of specific activity, fibrinolytic properties in human plasma in vitro and pharmacokinetic properties following bolus  
30 injection in hamsters.

**2. Reagents and Methods**

          The source of all reagents used in the present study has previously been reported (22), or is specified  
35 below. The template vector for mutagenesis, pMEX602sakB (i.e. pMEX.SakSTAR), has been described elsewhere (23). Restriction and modification enzymes were purchased from New England Biolabs (Leusden, The Netherlands),

00T260"064T0960

Boehringer Mannheim (Mannheim, Germany) or Pharmacia (Uppsala, Sweden). The enzymatic reactions were performed according to the supplier recommendation. The mutagenic oligonucleotides and primers were obtained from Eurogentec (Seraing, Belgium). Plasmid DNA was isolated using a purification kit from Qiagen (Hilden, Germany), as recommended. Transformation-competent *E. coli* cells were prepared by the well-known calcium phosphate procedure. Nucleotide sequence determination was performed on double strand plasmid DNA with the dideoxy chain termination method, using the T7 sequencing kit (Pharmacia, Uppsala, Sweden). Polymerase chain reactions (PCR) were performed using Taq polymerase from Boehringer Mannheim (Mannheim, Germany). The recombinant DNA methods required to construct the variants described in this example are well established (22, 27).

### 3. Construction of expression plasmids

The variants SakSTAR(K102C) and SakSTAR(K109C), were constructed by the spliced overlap extension polymerase chain reaction (SOE-PCR) (24) using pMEX.SakSTAR encoding SakSTAR as template. Two fragments were amplified by PCR (30 cycles: 1 sec at 94°C, 1 sec at 50°C, 10 sec at 72°C), the first one starting from the 5' end (primer 818A) of the staphylokinase gene to the region to be mutagenized (forward primer), the second one from this same region (backward primer) to the 3' end of the gene with primer 818D (5' CAAACAGCCAAGCTTCATT-CATTCAGC). The forward and backward primers shared an overlap of around 24 bp (for the construction of K102C: TAT GAT AAG AAT TGC AAA AAA GAA GAA (backward) and TTC TTC TTT TTT GCA ATT CTT ATC ATA (forward), for the construction of K109C: AAA AAG AAG AAA CGT GCT CTT TCC CTA (backward) and TAG GGA AAG AGC ACG TTT CTT CTT TTT (forward)). The two purified fragments were then assembled together in a second PCR reaction with the external primers 818A and 818D (30 cycles: 1 sec at 94°C, 1 sec at 50°C, 10 sec at 72°C). The amplified product



from this final reaction was purified, digested with EcoRI and HindIII and ligated into the corresponding site of pMEX.SakSTAR. For each construction, the sequence of the variant was confirmed by sequencing the entire coding 5 region.

#### 4. Expression and purification of SakSTAR variants

The SakSTAR variants were expressed and purified, as described below, from transformed *E. coli* grown in terrific broth (TB) medium (28). A 2 to 4 mL aliquot of an overnight saturated culture in LB medium was used to inoculate a 1 to 2 L culture in terrific broth supplemented with 100 µg/mL ampicillin. The culture was incubated with vigorous aeration and at 30°C. After about 16 hours incubation, IPTG (200 µmol/L) was added to the culture to induce expression from the *tac* promoter. After 3 hours induction, the cells were pelleted by centrifugation at 4,000 rpm for 20 min, resuspended in 1/10 volume of 0.01 mol/L phosphate buffer pH 6-6.5 and disrupted by sonication at 0°C. The suspension was centrifuged for 20 min at 20,000 rpm and the supernatant was stored at 4°C or at -20°C until purification. The material was purified essentially as described above (Example 2): cleared cell lysates containing the SakSTAR variants were subjected to chromatography on a 1.6 x 5 cm column of SP-Sephadex, followed by chromatography on a 1.6 x 8 cm column of phenyl-Sepharose. The SakSTAR containing fractions, localized by SDS-gel electrophoresis, were pooled for further analysis.

#### 5. Biochemical analysis

Protein concentrations were determined according to Bradford (29). SDS-PAGE was performed with the Phast System™ (Pharmacia, Uppsala, Sweden) using 10-15% gradient gels and Coomassie Brilliant blue staining, and the specific activities of SakSTAR solutions were determined with a chromogenic substrate assay carried out in microtiter plates (as described in

example 2). The specific activity of the different SakSTAR variants are summarized in Table 19.

Mutant SakSTAR(K102C) was essentially monomeric as visualized by SDS-PAGE and Coomassie Brilliant blue staining. Its specific activity was comparable to that of wild-type staphylokinase. In contrast, SakSTAR(K109C) showed a propensity to form dimers (> 60%). This resulted in a markedly increased specific activity in the plasminogen-coupled chromogenic substrate assay (see Table 19). Upon reduction with dithiothreitol (DTT) (20-fold molar excess during 1.5 hour at 37°C) and alkylation with iodoacetamide (100-fold molar excess during 1 hour at 37°C), the K109C dimer is converted into a stable monomer and its resulting specific activity is within the expected range towards wild-type staphylokinase (Table 19). This result confirms that formation of homodimers is the unique determinant for this large increase in specific activity. Dimeric SakSTAR(K109C) was separated from monomeric SakSTAR(K109C) by chromatography on Source S (Pharmacia) (5 x 50mm). Loading buffer was 10 mM phosphate, pH 6.0 and dimeric SakSTAR(K109C) was eluted by a salt gradient (up to 1 M) in the same buffer. The dimeric SakSTAR(K109C) (>95% pure) containing fractions, localized by SDS-gel electrophoresis, were pooled for further analysis.

#### 6. Chemical crosslinking of cysteine mutants of SakSTAR with polyethylene glycol

The thiol group of the cysteine mutant SakSTAR(K102C) was targeted for coupling with an activated polyethylene glycol, OPSS-PEG (Shearwater Polymers Europe, Enschede, The Netherlands). OPSS-PEG is a 5 kDa PEG molecule carrying a single activated thiol group at one end that react specifically at slightly alkaline pH with free thiols. Modification of SakSTAR(K102C) was achieved by incubating the molecule (100 µM) with a three-fold excess of SS-PEG in a 5 mM

phosphate, pH 7.9 solution at room temperature. The extent of the reaction was monitored by following the release of 2-thiopyridone from OPSS-PEG at 412 nm. After reaction (about 15 min), the excess of OPSS-PEG was removed by purifying the derivatized SakSTAR(K102C-PEG) on a 1.6 x 5 cm column of SP-Sephadex as described above (see Example 2). The SakSTAR(K102C-PEG) containing fractions, localized by optical density at 280 nm, were pooled for further analysis. SDS-PAGE analysis and Coomassie blue staining confirmed that PEG crosslinking on SakSTAR(K102C) was quantitative. As shown in Table 19, the specific activity of the PEG-derivative was only marginally affected when compared to that of wild-type staphylokinase.

15

7. Fibrinolytic properties of SakSTAR variants in human plasma in vitro

The fibrinolytic and fibrinogenolytic properties of SakSTAR variants were determined as previously described. Dose- and time-dependent lysis of <sup>125</sup>I-fibrin labeled human plasma clots submerged in human plasma was obtained with four molecules: SakSTAR(K109C) as dimer and as monomer (after reduction and alkylation with iodoacetamide), the monomeric SakSTAR(K102C) and the PEG-derivatized SakSTAR(K102C). Spontaneous clot lysis during the experimental period was ≤5% (not shown). Equi-effective concentrations of test compound (causing 50% clot lysis in 2 hrs; C<sub>50</sub>), determined graphically from plots of clot lysis at 2 hrs versus the concentration of plasminogen activator (not shown), were comparable to that of SakSTAR, for monomeric SakSTAR(K109C) and SakSTAR(K102C) (Table 19). However, it was observed that the C<sub>50</sub> for clot lysis by dimeric SakSTAR(K109C) was only 0.12 µg/ml, which is approximately three-fold lower than for wild-type staphylokinase. In contrast, a C<sub>50</sub> of 0.60 µg/ml was measured for SakSTAR(K102C-PEG), which is only two-fold higher than for wild-type staphylokinase. Thus,

dimerization of SakSTAR via disulfide bridges or increasing the size of the molecule via PEG-derivatization does not preclude the fibrinolytic activity of staphylokinase. While a PEG-molecule appears to reduce the diffusion and therefore fibrinolytic potency of the derivatized staphylokinase within a fibrin clot, dimerization of staphylokinase results in a synergistic fibrinolytic effect on human fibrin clots.

10 8. Pharmacokinetic properties of dimeric SakSTAR(K109C) and SakSTAR(K102C-PEG) following bolus injection in hamsters

The pharmacokinetic parameters of the disposition of dimeric SakSTAR(K109C) and SakSTAR(K102C-PEG) from blood were evaluated in groups of 4 hamsters following intravenous bolus injection of 100  $\mu\text{g/kg}$  SakSTAR variant. SakSTAR-related antigen was assayed using the ELISA described elsewhere. The ELISA was calibrated against each of the SakSTAR variants to be quantitated. Pharmacokinetic parameters included: initial half-life (in min),  $t_{1/2\alpha} = \ln 2/\alpha$ ; terminal half-life (in min),  $t_{1/2\beta} = \ln 2/\beta$ ; volume of the central (plasma) compartment (in mL),  $VC = \text{dose}/(A+B)$ ; area under the curve (in  $\mu\text{g}\cdot\text{min}\cdot\text{mL}^{-1}$ ),  $AUC = A/\alpha + B/\beta$ ; and plasma clearance (in mL $\cdot\text{min}^{-1}$ ),  $Cl_p = \text{dose}/AUC$  (32).

The disposition rate of staphylokinase-related antigen from blood following bolus injection of 100  $\mu\text{g/kg}$  of the selected SakSTAR variants in groups of 4 hamsters could adequately be described by a sum of two exponential terms by graphical curve peeling (results not shown), from which the pharmacokinetic parameters  $t_{1/2\alpha}$  and  $Cl_p$ , summarized in Table 19 were derived. The pharmacokinetic parameters of dimeric SakSTAR(K109C) and SakSTAR-(K102C-PEG) were markedly different from those of wild type SakSTAR. Initial plasma half-lives ( $t_{1/2(\alpha)}$ ) were 3.6 and 3.0 min and plasma clearances ( $Cl_p$ ) were 0.52 and 0.32 mL/min, for dimeric SakSTAR(K109C) and SakSTAR-(K102C-PEG), respectively. These results may be due to

09501490-092400  
007260-06410960

the increase of the Stokes radius of SakSTAR as a result of the dimerization or crosslinking with PEG. According to size-exclusion chromatography on Superdex50 by HPLC, dimeric SakSTAR(K109C) and SakSTAR(K102C-PEG) have  
5 apparent molecular weights of 33 kDa and 40 kDa, respectively.

#### EXAMPLE 14

Construction, purification and characterization of  
10 cysteine-substitution mutants of variants of  
staphylokinase with reduced immunogenicity

##### 1. Introduction

Based on the results of example 13, additional polyethylene glycol derivatives of SakSTAR variants were  
15 constructed, purified and characterized. The least immunogenic variants SakSTAR(E65D,K74R,E80A,D82A,K130T,K135R), (SY19), and SakSTAR(K35A,E65Q,K74Q,D82A,S84A,T90A,E99D,T101S,E108A,K109A,K130T,K135R,K136A,V137K), (SY141), were used as templates,  
20 with the proviso that the COOH-terminus of the latter was reverted to the wild type sequence, S84A was replaced with E80 and K74Q replaced with K74R, yielding SakSTAR(K35A,E65Q,K74R,E80A,D82A,T90A,E99D,T101S,E108A,K109A,K130T,K135R), (SY161). The introduced cysteine,  
25 which functions as acceptor of the polyethylene glycol molecule was located in the amino terminal region (preferably, but not exclusively, the Ser in position number 3 of the mature staphylokinase variant) in order to be released upon activation of staphylokinase (release  
30 of the 10 NH<sub>2</sub>-terminal amino acids); finally polyethylene glycol molecules of different molecular weights ( $M_n$  5,000 to 20,000) were used, substituted with either OPSS or maleimide.

The mutants described under this example are  
35 listed in Table 20. These variants were expressed in E.coli, purified and characterized in terms of specific activity, fibrinolytic properties in human plasma in vitro, pharmacokinetic properties following bolus

001260 06410950

injection in hamsters, thrombolytic properties following bolus injection in a hamster pulmonary embolism model, and absorption of antibodies from pooled immunized patient plasma (Pool 40).

5

## 2. Reagents and Methods

The source of all reagents used in the present study has previously been reported (22), or is specified below. The template vector for mutagenesis, pMEX602sakB (i.e. pMEX.SakSTAR), has been described elsewhere (23). Restriction and modification enzymes were purchased from New England Biolabs (Leusden, The Netherlands), Boehringer Mannheim (Mannheim, Germany) or Pharmacia (Uppsala, Sweden). The enzymatic reactions were performed according to the supplier recommendation. The mutagenic oligonucleotides and primers were obtained from Eurogentec (Seraing, Belgium). Plasmid DNA was isolated using a purification kit from Qiagen (Hilden, Germany), as recommended. Transformation-competent *E. coli* cells were prepared by the well-known calcium phosphate procedure. Nucleotide sequence determination was performed on double strand plasmid DNA with the dideoxy chain termination method, using the T7 sequencing kit (Pharmacia, Uppsala, Sweden). Polymerase chain reactions (PCR) were performed using Taq polymerase from Boehringer Mannheim (Mannheim, Germany). The recombinant DNA methods required to construct the variants described in this example are well established (22, 27).

## 3. Construction of expression plasmids

The variants SakSTAR(S3C,E65D,K74R,E80A,D82A,K130T,K135R), (SY19(S3C)), SakSTAR(S2C,S3C,E65D,K74R,E80A,D82A,K130T,K135R), (SY19(2SC,3SC)), SakSTAR(S3C,K35A,E65Q,K74Q,D82A,S84A,T90A,E99D,T101S,E108A,K109A,K130T,K135R,K136A,V137K), (SY141(S3C)), SakSTAR(S2C,S3C,K35A,E65Q,K74Q,D82A,S84A,T90A,E99D,T101S,E108A,-K109A,K130T,K135R,K136A,V137K), (SY141(S2C,S3C)), SakSTAR(S3C,K35A,E65Q,K74Q,E80A,D82A,T90A,E99D,T101S,E108A,

K109A, K130T, K135R), (SY160(S3C)) and SakSTAR(S3C, K35A, E65Q, K74R, E80A, D82A, T90A, E99D, T101S, E108A, K109A, K130T, K135R), (SY161(S3C)), were constructed by the spliced overlap extension polymerase chain reaction (SOE-PCR) (24) using pMEX.SakSTAR encoding SakSTAR as template, two fragments were amplified by PCR (30 cycles: 1 sec at 94°C, 1 sec at 50°C, 10 sec at 72°C), the first one starting from the 5' end (primer 818A) of the staphylokinase gene to the region to be mutagenized (forward primer), the second one from this same region (backward primer) to the 3' end of the gene with primer 818D (5' CAAACAGCCAAGCTTCATTCATTCAGC). The forward and backward primers shared an overlap of around 24 bp. The two purified fragments were then assembled together in a second PCR reaction with the external primers 818A and 818D (30 cycles: 1 sec at 94°C, 1 sec at 50°C, 10 sec at 72°C). The amplified product from this final reaction was purified, digested with EcoRI and HindIII and ligated into the corresponding site of pMEX.SakSTAR. For each construction, the sequence of the variant was confirmed by sequencing the entire SakSTAR coding region.

#### 4. Expression and purification of SakSTAR variants

The SakSTAR variants were expressed and purified, as described below, from transformed *E. coli* grown in terrific broth (TB) medium (28). A 2 to 4 mL aliquot of an overnight saturated culture in LB medium was used to inoculate a 1 to 2 L culture in terrific broth supplemented with 100 µg/mL ampicillin. The culture was incubated with vigorous aeration and at 30°C. After about 16 hours incubation, IPTG (200 µmol/L) was added to the culture to induce expression from the tac promoter. After 3 hours induction, the cells were pelleted by centrifugation at 4,000 rpm for 20 min, resuspended in 1/10 volume of 0.01 mol/L phosphate buffer pH 6-6.5 and disrupted by sonication at 0°C. The suspension was centrifuged for 20 min at 20,000 rpm and the supernatant was stored at 4°C or at -20°C until purification. The

007260 0640960

material was purified essentially as described above (Example 2): cleared cell lysates containing the SakSTAR variants were subjected to chromatography on a 1.6 x 5 cm column of SP-Sephadex, followed by chromatography on a 1.6 x 8 cm column of phenyl-Sepharose. The SakSTAR containing fractions, localized by SDS-gel electrophoresis, were pooled for further analysis.

#### 5. Biochemical analysis

Protein concentrations were determined according to Bradford (29). SDS-PAGE was performed with the Phast System™ (Pharmacia, Uppsala, Sweden) using 10-15% gradient gels and Coomassie Brilliant blue staining, and the specific activities of SakSTAR solutions were determined with a chromogenic substrate assay carried out in microtiter plates (as described in example 2).

#### 6. Chemical crosslinking of cysteine mutants of SakSTAR with polyethylene glycol

The thiol group of the cysteine mutants was targeted for coupling with an activated polyethylene glycol, either OPSS-PEG or MAL-PEG (Shearwater Polymers Europe, Enschede, The Netherlands). OPSS-PEG is a 5 kDa PEG molecule carrying a single activated thiol group at one end that reacts specifically at slightly alkaline pH with free thiols. MAL-PEG is a 5 kDa, 10 kDa or 20 kDa molecule carrying a maleimide group that reacts specifically with thiol groups under mild conditions in the presence of other functional groups. Modification of the variants was achieved by incubating the molecule (100 μM) with a three-fold excess of OPSS-PEG or MAL-PEG in a 5 mM phosphate, pH 7.9 solution at room temperature. After reaction (about 15 min), the excess of OPSS-PEG or MAL-PEG was removed by purifying the derivatized SakSTAR variant on a 1.6 x 5 cm column of SP-Sephadex as described above (see Example 2). The "pegylated" SakSTAR variant containing fractions, localized by optical densi-



ty at 280 nm, were pooled for further analysis. SDS-PAGE analysis and Coomassie blue staining confirmed that PEG crosslinking was quantitative. As shown in Table 20, the specific activities of the PEG-derivatives were only marginally affected when compared to that of wild-type staphylokinase.

7. Fibrinolytic properties of SakSTAR variants in human plasma in vitro

10           The fibrinolytic and fibrinogenolytic properties of SakSTAR variants were determined as previously described. Dose- and time-dependent lysis of  $^{125}\text{I}$ -fibrin labeled human plasma clots submerged in human plasma was obtained with all molecules tested.

15 Equi-effective concentrations of test compound (causing 50% clot lysis in 2 hrs;  $C_{50}$ ), determined graphically from plots of clot lysis at 2 hrs versus the concentration of plasminogen activator (not shown), were comparable to or only slightly lower than that of SakSTAR (Table 20). The

20  $C_{50}$  for clot lysis by variants derivatized with P20 (PEG with  $M_r$  20 kDa) was about twice as high as the non-derivatized variants. Thus increasing the size of the molecule via PEG-derivatization does not markedly affect the fibrinolytic activity of staphylokinase. The

25 PEG-molecules appear to reduce the diffusion and therefore fibrinolytic potency of the derivatized staphylokinase within a fibrin clot, but this appears to be less pronounced with variants substituted in their  $\text{NH}_2$ -terminal region - which is released during processing

30 of staphylokinase - than with variants substituted in the core of the molecule (cfr. Tables 19 and 20).

8. Pharmacokinetic properties of SakSTAR variants chemically modified with polyethylene glycol following bolus injection in hamsters

35           The pharmacokinetic parameters of the disposition of the pegylated variants from blood were evaluated in groups of 4 hamsters following intravenous

bolus injection of 100  $\mu\text{g/kg}$  SakSTAR variant.

SakSTAR-related antigen was assayed using the ELISA described elsewhere. The ELISA was calibrated against each of the SakSTAR variants to be quantitated.

- 5 Pharmacokinetic parameters included: initial half-life (in min),  $t_{1/2\alpha} = \ln 2/\alpha$ ; terminal half-life (in min),  $t_{1/2\beta} = \ln 2/\beta$ ; volume of the central (plasma) compartment (in mL),  $VC = \text{dose}/(A+B)$ ; area under the curve (in  $\mu\text{g}\cdot\text{min}\cdot\text{mL}^{-1}$ ),  $AUC = A/\alpha + B/\beta$ ; and plasma clearance (in mL $\cdot\text{min}^{-1}$ ),  $Clp = \text{dose}/AUC$  (32).

The disposition rate of staphylokinase-related antigen from blood following bolus injection of 100  $\mu\text{g/kg}$  of the selected SakSTAR variants in groups of 4 hamsters could adequately be described by a sum of two exponential terms by graphical curve peeling (results not shown), from which the plasma clearances  $Clp$ , summarized in Table 20 were derived. The clearances of pegylated variants were markedly different from those of wild type SakSTAR and were inversely proportional to the molecular weight of the PEG molecules, with an average reduction of 5-fold with PEG 5 kDa, 10-fold with PEG 10 kDa and 30-fold with PEG 20 kDa. These results may be due to the increase of the Stokes radius of SakSTAR as a result of crosslinking with PEG.

25

#### EXAMPLE 15

Comparative thrombolytic efficacy and clearance of SakSTAR(S3C-P20,E65D,K74R, E80A,D82A,K130T,K135R), (SY19(S3C-P20)), in two patients with acute myocardial infarction

30

Large scale purification and conditioning of the SakSTAR variant for use in vivo

Material was purified to homogeneity out of culture volumes of 18 liters. The endotoxin content was below 1 IU/mg. Gel filtration on HPLC revealed a single main symmetrical peak in the chromatographic range of the column, representing >98% of the eluted material (total area under the curve) (not shown). SDS gel

007260-06470560

electrophoresis of a 30  $\mu$ g sample revealed single main component. The preparation sterilized by filtration proved to be sterile on 3 day testing as described in methods. Intravenous bolus injection of the SakSTAR 5 variant in 5 mice (3 mg/kg body weight), did not provoke any acute reaction, nor reduced weight gain within 8 days, in comparison with mice given an equal amount of saline (not shown).

Two patients with acute myocardial infarction 10 were given a bolus injection of 5 mg SY19(S3C-P20). These patients had a complete recanalization of the occluded infarct-related artery as determined by coronary angiography at 90 min after the bolus injection. The material was cleared from the plasma with an initial 15 half-life of 3 to 4 hours, as compared to 4 to 6 minutes for wild-type SakSTAR. These data confirm that pegylated variants of SakSTAR may be useful for thrombolytic therapy by single bolus injection at a reduced dose.

## 20 CONCLUSION

In summary, the present invention shows that staphylokinase variants with markedly reduced antibody induction but intact thrombolytic potency can be generated. This observation constitutes the first case in 25 which a heterologous protein, with the use of protein engineering techniques, is rendered significantly less immunogenic in man without reducing its biological activity. In addition, the present invention shows that selective chemical modification of staphylokinase or its 30 variants with polyethylene glycol of varying molecular weights is feasible, resulting in a reduction of the plasma clearance proportional to the molecular weight. In the preferred embodiment an amino acid in the NH<sub>2</sub>-terminal region of staphylokinase, the portion that is removed by 35 processing, is substituted with Cys and the introduced thiol group is chemically modified with OPSS-PEG or MAL-PEG. This results in homogeneous products which, upon single intravenous bolus injection in experimental

animals and in patients have a maintained thrombolytic potency at markedly reduced doses.

1997-1998		1998-1999		1999-2000		2000-2001		2001-2002		2002-2003		2003-2004		2004-2005		2005-2006		2006-2007		2007-2008		2008-2009		2009-2010		2010-2011		2011-2012		2012-2013		2013-2014		2014-2015		2015-2016		2016-2017		2017-2018		2018-2019		2019-2020		2020-2021		2021-2022		2022-2023		2023-2024		2024-2025		2025-2026		2026-2027		2027-2028		2028-2029		2029-2030		2030-2031		2031-2032		2032-2033		2033-2034		2034-2035		2035-2036		2036-2037		2037-2038		2038-2039		2039-2040		2040-2041		2041-2042		2042-2043		2043-2044		2044-2045		2045-2046		2046-2047		2047-2048		2048-2049		2049-2050		2050-2051		2051-2052		2052-2053		2053-2054		2054-2055		2055-2056		2056-2057		2057-2058		2058-2059		2059-2060		2060-2061		2061-2062		2062-2063		2063-2064		2064-2065		2065-2066		2066-2067		2067-2068		2068-2069		2069-2070		2070-2071		2071-2072		2072-2073		2073-2074		2074-2075		2075-2076		2076-2077		2077-2078		2078-2079		2079-2080		2080-2081		2081-2082		2082-2083		2083-2084		2084-2085		2085-2086		2086-2087		2087-2088		2088-2089		2089-2090		2090-2091		2091-2092		2092-2093		2093-2094		2094-2095		2095-2096		2096-2097		2097-2098		2098-2099		2099-2100		2100-2101		2101-2102		2102-2103		2103-2104		2104-2105		2105-2106		2106-2107		2107-2108		2108-2109		2109-2110		2110-2111		2111-2112		2112-2113		2113-2114		2114-2115		2115-2116		2116-2117		2117-2118		2118-2119		2119-2120		2120-2121		2121-2122		2122-2123		2123-2124		2124-2125		2125-2126		2126-2127		2127-2128		2128-2129		2129-2130		2130-2131		2131-2132		2132-2133		2133-2134		2134-2135		2135-2136		2136-2137		2137-2138		2138-2139		2139-2140		2140-2141		2141-2142		2142-2143		2143-2144		2144-2145		2145-2146		2146-2147		2147-2148		2148-2149		2149-2150		2150-2151		2151-2152		2152-2153		2153-2154		2154-2155		2155-2156		2156-2157		2157-2158		2158-2159		2159-2160		2160-2161		2161-2162		2162-2163		2163-2164		2164-2165		2165-2166		2166-2167		2167-2168		2168-2169		2169-2170		2170-2171		2171-2172		2172-2173		2173-2174		2174-2175		2175-2176		2176-2177		2177-2178		2178-2179		2179-2180		2180-2181		2181-2182		2182-2183		2183-2184		2184-2185		2185-2186		2186-2187		2187-2188		2188-2189		2189-2190		2190-2191		2191-2192		2192-2193		2193-2194		2194-2195		2195-2196		2196-2197		2197-2198		2198-2199		2199-2200		2200-2201		2201-2202		2202-2203		2203-2204		2204-2205		2205-2206		2206-2207		2207-2208		2208-2209		2209-2210		2210-2211		2211-2212		2212-2213		2213-2214		2214-2215		2215-2216		2216-2217		2217-2218		2218-2219		2219-2220		2220-2221		2221-2222		2222-2223		2223-2224	
-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--

**Table 1: Alanine-to-wild-type" reversal variants of "charged-cluster-to-alanine" mutants of SakSTAR: Association constants ( $K_A \times 10^7 \text{ mol/L}^{-1}$ ) for the binding to insolubilized murine monoclonal antibodies (MAbs), and absorption (percent) of antibodies of immunized patient plasma**

Variant	Exp. (mg/L)	Spec. Act. (kU/mg)	murine MAbs												SakSTAR patient plasma					
			Epitope I						Epitope II						Epitope III					
			17G11	25A2	30A2	2B12	3G10	18F12	14H5	28H4	32B2	7F10	7H11	23E1	40C8	24C4	1A10	Pool	Subpool B	Subpool C
SakSTAR	22	13	29	7.8	11	38	7.4	19	7.7	2.4	4.0	14	5.4	2.9	0.6	95	95	95		
SakSTAR(K35A.E38A)	97		15	22	42	11	7.9	110	10	15	12	2.2	<0.1	<0.1	<0.1	1.0	1.0	93	91	94
SakSTAR(K74A.E75A.R77A)	110		11	<0.1	<0.1	<0.1	<0.1	150	17	28	14	3.3	2.4	11	4.0	2.1	0.9	55	43	95
SakSTAR(K35A.E38A.K74A.E75A.R77A)	50		11	<0.1	<0.1	<0.1	<0.1	110	36	26	15	2.0	<0.1	<0.1	<0.1	1.5	1.2	52	41	92
SakSTAR(E38A.K74A.E75A.R77A)	43		11	<0.1	<0.2	<0.1	<0.1	140	39	26	15	2.1	<0.1	3.2	3.7	1.6	1.1	50	44	95
SakSTAR(K35A.K74A.E75A.R77A)	56		9.2	<0.1	0.15	<0.1	<0.1	52	14	29	8.8	2.3	<0.1	1.8	<0.1	1.8	0.8	46	43	95
SakSTAR(K35A.E38A.E75A.R77A)	44		11	0.3	0.1	0.2	<0.1	75	9.8	12	7.3	1.6	<0.1	<0.1	<0.1	0.53	0.64	92	87	94
SakSTAR(K35A.E38A.K74A.R77A)	41		8.8	2.9	<0.1	2.0	0.33	110	29	31	10	2.0	<0.1	<0.1	<0.1	0.63	0.74	56	50	93
SakSTAR(K35A.E38A.K74A.E75A)	19		13	<0.1	0.1	<0.1	<0.1	180	41	37	15	1.6	<0.1	<0.1	<0.1	1.2	0.45	48	41	92
SakSTAR(E38A.E75A.R77A)	88		11	0.6	0.15	0.4	0.3	79	12	15	10	2.0	<0.1	2.6	4.7	1.1	0.81	95	88	95
SakSTAR(E38A.E75A)	66		16	0.3	<0.1	<0.1	0.9	56	11	13	8.9	2.0	<0.1	20	4.8	1.3	1.6	91	90	95
SakSTAR(K35A.E75A.R77A)	68		9.2	<0.1	<0.1	<0.1	<0.1	60	7.0	13	11	3.3	<0.1	1.5	<0.1	0.8	1.1	88	89	95
SakSTAR(K35A.E75A)	150		17	0.12	<0.1	0.16	0.14	40	7.2	13	9.2	4.2	<0.1	1.8	<0.1	1.4	1.5	94	93	95
SakSTAR(K74A)	100		12	7.6	0.17	4.4	2.1	55	15	33	14	3.6	2.9	14	4.9	3.4	1.2	59	45	95
SakSTAR(E75A)	140		13	1.2	<0.1	<0.1	<0.1	46	8.5	14	12	3.4	4.5	18	5.0	1.2	2.1	95	93	95
SakSTAR(K74A.E75A.R77A.E80A.D82A)	50		14	<0.1	<0.1	<0.1	<0.1	180	19	33	19	3.7	<0.1	<0.1	<0.1	<0.1	1.2	49	29	89
SakSTAR(E80A.D82A)	130		7.3	12	2.1	6.5	5.9	79	6.1	8.4	7.8	1.9	<0.1	<0.1	<0.1	<0.1	0.44	89	83	92
SakSTAR(E80A)	160		13	13	3.3	7.9	10	35	7.4	17	8.6	2.1	<0.1	16	3.6	<0.1	1.7	94	93	95
SakSTAR(D82A)	160		17	12	4.8	7.3	11	31	7.8	17	12	2.7	<0.1	0.18	<0.1	<0.1	2.3	95	93	95
SakSTAR(E75A.D82A)	170		20	15	3.1	6.6	7.2	69	8.1	15	14	4.9	0.17	0.7	0.5	0.1	1.4	95	95	95

Apparent association constants  $\geq 10$ -fold lower than those of wild-type SakSTAR are represented in bold type; Spec. Act.  $\geq 100,000 \text{ HU/mg}$  represented in bold type;  $\leq 60\%$  absorption represented in bold type.

Table 2: Baseline characteristics and treatment outcome of the patients with peripheral arterial occlusion treated with SakSTAR, SakSTAR(K74A) or SakSTAR(K74A,E75A,R77A)

Compound Patient Id.	Gender	Age (yrs)	Clinical ischemia	Locus of occlusion	Age of occlusion (days)	Length of occlusion (cm)	Recanalization by thrombolysis	Total dose of thrombolytic agent (mg)	Total duration of infusion (hrs)	Additional therapy
<b>SakSTAR</b>										
MEE	F	67	Rest pain	Left SFA	30	8	complete	7.0	5.0	PTA
FOR	M	68	Claudication	Left IA (stent)	14	18	complete	6.5	4.5	PTA + stent
DAN	M	73	Claudication	Right SFA	30	6	complete	7.5	5.5	PTA
BER	F	63	Rest pain	Left FT graft	18	55	complete	18	28	PTA
DAM	F	43	Acute	Left brachial and radial artery	2	7	complete	19	17	PTA + stent
TOR	M	68	Claudication	Right SFA (popliteal aneurysm)	50	12	complete	6.0	4.0	PTA + femoropopliteal bypass graft
CLA	M	74	Acute	Left PA	1.5	20	complete	9.0	7.0	-
MAN	M	65	Acute	Left EIA (stent)	4	20	complete	6.5	4.5	(amputation left digit V)
MAT	M	64	Subacute	Right FP graft	3	45	complete	8.0	6.0	(-)
Mean $\pm$ SEM		65 $\pm$ 3.0			17 $\pm$ 5.6	21 $\pm$ 5.8		9.7 $\pm$ 1.7	9.1 $\pm$ 2.7	
<b>SakSTAR(K74A)</b>										
LIE	M	70	Subacute	Right FF graft	10	48	complete	11	9.0	PTA
ENG	M	50	Claudication	Right SFA	28	10	complete	12	10	PTA
COX	F	48	Claudication	Right PA graft	25	7	partial	15	15	PTA
MAN	F	68	Claudication	Right SFA	$\geq 120$	9	complete	9.0	7.0	PTA
VHE	M	47	Acute	Right IF graft	10	54	complete	18	16	Surgical graft revision
MUL	F	51	Acute	Right IF and FP graft	1	63	complete	16	20	PTA
BUR	F	67	Rest pain	Right TF trunc	9.0	38	partial	18	21	-
NIJ	F	60	Rest pain	Left AF graft	23	78	complete	15	21	-
POE*	M	49	Subacute	Right TF trunc	2	30	partial	6.0	4.0	rt-PA, surgical graft lengthening
VBE	M	39	Subacute	Right BA (embolism)	20	28	complete	18	23	Stent right SC artery, first rib resection
SME	F	50	Subacute	TF trunc	18	32	complete	21	19	None
WOL	M	67	Subacute	Right PA	4	25	complete	16	22	-
Mean $\pm$ SEM		56 $\pm$ 3.0			23 $\pm$ 9.2	35 $\pm$ 6.4		15 $\pm$ 1.2	16 $\pm$ 1.9	
<b>SakSTAR(K74A,E75A,R77A)</b>										
JAC	F	65	Acute	Right BA and UA	0.3	5	complete	14	12	-
MAE	M	74	Rest pain	Left SFA	10	50	complete	9.0	7.0	PTA
CRA	F	52	Claudication	Right IA and FA artery	14	28	complete	25	23	PTA + stent
VDB	M	68	Claudication	Left SFA	90	12	complete	9.0	7.0	PTA
DUN	M	71	Subacute	Left SFA	14	6	complete	9.0	7.0	PTA
DEL	M	59	Acute	Right FT graft	3	42	complete	9.0	7.0	PTA
Mean $\pm$ SEM		63 $\pm$ 3.3			22 $\pm$ 14	24 $\pm$ 7.8		13 $\pm$ 2.6	11 $\pm$ 2.6	

AF: aortofemoral; BA: brachial artery; CIA: common iliac artery; FF: femorofibular; FP: femoropopliteal; FT: femorotibial; IA: iliac artery; IF: iliofemoral; PA: popliteal artery; PTA: percutaneous transluminal angioplasty; SFA: superficial femoral artery; TF: tibiofibular; UA: ulnar artery. \*Previous treatment with SakSTAR in 1994

**Table 3: Alanine-substitution variants of SakSTAR: Association constants ( $K_A \times 10^7 \text{ mol/L}^{-1}$ ) for binding to insolubilized murine monoclonal antibodies (Mab) and absorption (percent) of antibodies of immunized patient plasma**

Variant		Exp (mg/L)	Spec Act (kU/mg)	murine MAbs										SakSTAR patient plasma									
				Epitope cluster I					Epitope cluster II					Pool	Subpool B	Subpool C							
				17G11	26A2	30A2	2812	3G10	18F12	14H5	28H4	32B2	7F10				7H11	25E1	40C8	24C4	7A10		
SakSTAR				120	93	13	29	78	11	38	74	19	77	24	40	14	54	29	0.6	95	95	95	
SakSTAR(S34G,G36R,H43R)				120	164	17	74	19	35	18	25	>18	>14	11	0.6	11	16	6.3	2.01	87	76	75	
SakSTAR(F4A)				150	113	21	12	92	97	12	23	17	10	04	13	39	08	45	3.81	95	95	95	
SakSTAR(D5A,K6A)				24	118	16	51	29	15	22	16	26	18	10	0.93	11	1.1	18	0.751	95	95	95	
SakSTAR(K8A,K10A)				78	22	49	66	23	16	44	84	20	14	26	24	16	96	11	0.5	96	95	95	
SakSTAR(Y9A)				24																			
SakSTAR(K11A,D13A,D14A)				LE																			
SakSTAR(D13A)				6	46	24	61	20	37	34	11	19	4.5	4.4	87	1.5	24	11	3.6	<0.1	95	94	95
SakSTAR(D14A)				14	30	51	13	40	66	31	38	77	12	15	22	27	6.0	32	5.6	<0.1	95	94	95
SakSTAR(S16A)				41	160	81	16	45	84	90	15	61	13	21	3.6	40	80	4.3	2.5	0.5	95	95	95
SakSTAR(Y17A,F18A)				27	30	13	22	33	10	95	21	46	67	12	25	1.2	18	65	3.4	<0.1	95	95	95
SakSTAR(E19A,P20A)				16	9	11	19	33	92	12	15	61	11	16	10	1.1	15	51	33	<0.1	93	93	95
SakSTAR(T21A)				54	170	48	15	24	87	96	32	11	24	18	13	18	96	29	56	0.6	95	95	95
SakSTAR(P21A)				41	67	14	31	44	14	22	41	53	37	13	10	40	11	76	31	1.9	91	95	95
SakSTAR(Y24A)				10	40	17	33	43	13	11	33	43	70	12	40	04	14	68	48	<0.1	95	95	95
SakSTAR(L25A)				LE																			
SakSTAR(M26A)				LE																			
SakSTAR(V27A)				62	50	33	15	16	78	74	12	29	37	33	20	13	59	28	42	1.2	95	95	95
SakSTAR(N28A)				90	<5	58	29	21	70	55	27	10	20	25	21	21	56	21	21	07	95	95	95
SakSTAR(N28A,V29A)				32	45	18	30	25	20	18	20	14	20	24	27	33	20	11	50	20	93	95	95
SakSTAR(T30A)				52	140	74	13	21	70	61	76	34	51	13	33	56	12	34	54	08	94	95	95
SakSTAR(V32A)				78	45	10	96	24	62	78	56	17	12	14	14	<0.1	<0.1	<0.1	<0.1	2.2	90	93	95
SakSTAR(D13A,K15A)				130		121	19	14	14	19	15	24	32	10	53	14	51	38	51	301	95	95	95
SakSTAR(S34A)				29	110	17	24	46	95	11	28	11	22	15	29	31	88	38	20	02	95	95	93

Variant	Exp (mg/L)	Spec. Act (AU/mg)	murine MAbs												Epitope cluster III				SakSTAR patient plasma	
			Epitope cluster I				Epitope cluster II				Epitope cluster III				Pool	Subpool B	Subpool C			
			7B11 26A	30A7 28B2	3G10 28H4	3G10 28H4	18F12 14H5	28H4 28B2	7F10 28H4	7H11 25E1	40C8 24C4	1A10 24C4	91	88						
SakSTAR(K35A)	230	6.4	14	33	80	7.4	31	11	12	11	2.6	<0.1	1.3	0.2	1.7	0.8	91	88	95	
SakSTAR(K35A.E38A)	97	15	22	42	11	7.9	110	10	15	12	2.2	<0.1	<0.1	<0.1	1.0	1.0	93	91	94	
SakSTAR(G36A)	14	72	3.5	9.8	1.5	6.5	52	42	17	9.2	1.4	<0.1	<0.1	0.9	5.0	1.0	86	83	78	
SakSTAR(N37A)	40	110	5.6	11	3.0	10	11	20	41	14	10	1.4	5.3	3.5	3.6	0.8	95	95	95	
SakSTAR(L39A.L40A)	8	<5	14	15	3.1	5.1	8.0	27	16	6.3	12	1.2	5.4	3.2	2.1	0.9	93	93	95	
SakSTAR(S41A.P42A)	24	48	10	25	4.1	13	12	11	3.0	1.9	27	3.2	1.5	4.8	3.6	1.1	95	95	95	
SakSTAR(H43A)	37	69	15	28	9.7	18	7.6	<0.1	<0.1	<0.1	9.1	1.5	2.0	2.3	7.8	1.6	95	95	95	
SakSTAR(H43A.Y44A)	13	<5	10	22	3.7	17	15	<0.1	<0.1	<0.1	1.4	3.0	2.3	1.1	5.2	0.1	95	95	95	
SakSTAR(V45A)	19	<5	16	5.6	1.4	4.8	6.3	2.2	0.2	1.7	32	2.6	2.1	8.3	1.1	1.6	91	92	95	
SakSTAR(E46A.K50A)	LE																			
SakSTAR(F47A)	6	<5	<0.1	4.0	1.0	3.9	3.4	5.7	2.7	2.8	8.5	1.7	0.9	8.8	3.0	3.0	90	82	97	
SakSTAR(I49A)	2	43	2.7	2.7	7.8	23	22	35	4.4	11	6.2	1.1	2.0	5.7	2.0	1.7	95	95	95	
SakSTAR(K50A)	15	42	<0.1	13	2.9	7.8	8.7	4.6	8.3	12	2.2	0.5	2.8	6.4	4.0	2.3	95	94	95	
SakSTAR(T53A.T54A)	14	68	0.9	19	2.7	7.6	7.8	4.1	6.7	12	1.5	1.5	1.9	5.1	2.3	1.0	93	94	95	
SakSTAR(L55A)	LE																			
SakSTAR(T56A)	17	150	5.5	15	3.2	12	13	5.6	5.3	12	11	2.0	3.5	6.1	2.7	4.3	94	92	95	
SakSTAR(K57A.E58A.K59A)	94	94	4	8.7	6.0	7.3	27	16	14	6.7	5.6	0.52	0.36	1.7	0.42	1.0				
SakSTAR(I60A)	11	96	12	20	2.9	11	13	27	6.4	2.5	2.7	1.5	0.7	5.8	2.9	1.7	95	95	95	
SakSTAR(E61A.E65A)	80		9.5	>10	8.8	21	29	>11	>16	6.6	>7.2	4.6	0.5	4.6	2.0	5.9				
SakSTAR(Y62A.Y61A)	24	<5	<0.1	4.3	0.3	2.1	1.9	11	2.2	3.1	8.4	1.7	0.6	9.2	3.6	3.8	89	81	95	
SakSTAR(Y61A)	4	<5	<0.1	18	3.7	9.6	13	17	5.3	3.3	1.5	1.1	2.2	5.3	4.3	1.0	89			

\_\_\_\_\_



**Alanine-substitution variants of SakSTAR: Association constants ( $K_A \times 10^4 \text{ mol/L}^{-1}$ ) for binding to insolubilized murine monoclonal antibodies (Mab) and absorption (percent) of antibodies of immunized patient plasma**

Virus	Exp (mg/L)	Spec Act (iU/mg)	murine MAbs										SalSTAR patient plasma						
			Epitope cluster I			Epitope cluster II				Epitope cluster III			Pool	Subpool B	Subpool C				
			17G11	26A2	30A2	2B12	3G10	10F12	14H5	28H4	7F10	7H11	25E1	28C4	1A10				
SakSTAR(Y71A)	20	<3	9.5	<0.1	<0.1	<0.1	4.8	3.3	7.3	11	8.9	2.0	0.5	9.0	2.5	1.4	63	44	93
SakSTAR(Y17A,K74A)	24	<5	18	<0.1	<0.1	<0.1	<0.1	19	6.7	23	9.9	3.2	2.7	13	4.0	1.6	47	28	87
SakSTAR(K74A)	80	69	4.4	2.7	0.2	2.2	1.1	17	5.2	14	7.6	2.2	2.0	6.8	3.3	1.8	64	58	95
SakSTAR(K74A,E75A,R77A)		68	9.2	<0.1	<0.1	<0.1	<0.1	60	7.0	13	11	3.3	<0.1	1.5	<0.1	0.8	88	89	95
SakSTAR(K74A,R77A)	34	41	3.5	1.8	0.2	1.5	0.4	20	2.4	10	2.1	1.8	1.7	2.3	2.2	1.2	74	60	95
SakSTAR(E75A)		140	13	1.2	<0.1	<0.1	<0.1	46	8.5	14	12	3.4	4.5	18	5.0	1.2	95	93	95
SakSTAR(F76A)	9	90	20	9.6	1.0	2.7	3.9	13	6.2	20	15	1.7	0.3	5.9	2.1	1.2	94	92	95
SakSTAR(V78A,V79A)	23	68	12	23	4.0	10	17	21	18	34	28	2.3	1.6	4.7	<0.1	0.5	93	93	95
SakSTAR(E80A)		160	13	13	3.1	7.9	10	15	7.4	17	8.6	2.1	<0.1	1.6	3.6	<0.1	94	93	95
SakSTAR(E80A,D82A)		130	7.3	12	2.1	6.5	5.9	79	6.1	8.4	7.8	1.9	<0.1	<0.1	<0.1	0.4	89	83	92
SakSTAR(L81A)	23	28	12	33	1.6	40	11	52	11	17	17	3.9	1.4	5.2	7.1	4.6	88	95	95
SakSTAR(D82A)		160	17	12	4.8	7.3	11	31	7.8	17	12	2.7	<0.1	0.2	<0.1	<0.1	95	93	95
SakSTAR(D82A,S84A)	72	130	8.3	14	2.6	8.1	8.5	23	3.8	12	11	1.7	<0.1	<0.1	1.4	<0.1	91	91	95
SakSTAR(S84A)	12/26	89	8.0	16	3.8	8.6	10	90	8.3	11	36	1.8	2.2	1.6	3.0	3.5	95	95	95
SakSTAR(K86A,E88A)		73	17.2	1.4	3.7	6.0	4.6	5.7	4.9	7.7	15	4.4	<0.1	5.4	0.80	1.9			
SakSTAR(I87A)	18	98	6.7	21	2.8	8.6	9.1	10	3.6	11	7.4	2.7	1.1	7.8	3.4	4.5	95	95	95
SakSTAR(V89A)	20	87	4.6	11	2.6	6.6	2.2	28	7.2	7.3	3.0	1.3	1.2	5.1	2.9	3.1	95	95	95
SakSTAR(T90A)	78	120	6.0	12	0.9	3.7	3.1	20	4.8	7.2	<0.1	<0.1	2.1	6.6	2.6	2.1	95	95	95
SakSTAR(Y91A)	5	53	6.0	16	3.0	7.0	13	28	8.2	16	0.6	2.1	1.4	3.7	1.6	1.6	95	95	95
SakSTAR(Y92A)	16	120	1.6	23	4.1	1.3	12	29	7.3	18	<0.1	1.7	4.4	10	3.9	5.9	95	95	95
SakSTAR(E91A,K94A)		97	18.2	19	1.3	30	24	18	11	>10	90	0.88	1.4	11	2.4	7.0			
SakSTAR(K94A,N95A,K97A)	12	8	NT														95	94	95
SakSTAR(N95A)	25	360	10	18	4.0	10	11	50	13	14	4.9	2.3	3.7	7.3	4.7	2.9	95	94	95
SakSTAR(K96A,K97A,K98A)		47	12.8	41	23	37	90	>16	91	19	16	0.41	0.58	17	12	13			
SakSTAR(E99A)	24	42	7.4	15	4.0	8.4	8.9	22	2.7	4.7	<0.1	<0.1	2.1	6.2	7.3	1.4	92	91	92
SakSTAR(E99A,E100A)	1E																		
SakSTAR(I101A)	21	85	4.6	11	2.1	6.6	7.3	30	2.8	16	0.7	1.0	1.3	5.4	2.4	2.9	95	95	95

**Table 3 - cont'd:** Alanine-substitution variants of SakSTAR: Association constants ( $K_A \times 10^7 \text{ mol/L}^{-1}$ ) for binding to insolubilized murine monoclonal antibodies (Mab) and absorption (percent) of antibodies of immunized patient plasma

Variant	Exp. (mg/L)	Spec. Act (kU/mg)	murine MAbs															SakSTAR patient plasma		
			Epitope cluster I					Epitope cluster II					Epitope cluster III					Pool	Subpool B	Subpool C
			7G11	26A2	30A2	2812	3G10	18F12	14R5	28H4	32B2	7F10	7H11	25E1	40C8	24C4	1A10			
SakSTAR(K102A)	12	89	49	12	37	65	63	30	31	15	82	61	08	42	19	10	06	95	93	95
SakSTAR(S103A)	67	210	90	16	50	94	91	19	59	13	13	36	39	83	47	28	09	94	95	95
SakSTAR(F104A)	14	55	58	19	48	14	27	71	50	14	48	<0.1	0.4	7.6	3.4	11	1.3	95	93	95
SakSTAR(I106A)	2	93	23	13	30	74	67	55	52	17	11	14	18	31	18	12	0.5	95	95	95
SakSTAR(T107A)	32	130	52	15	34	98	10	32	87	47	14	19	31	63	32	50	08	94	94	95
SakSTAR(E108A K109A)		170	116	51	72	19	51	28	15	21	21	12	0.43	69	14	10	19			
SakSTAR(F111A)	5	49	37	16	38	13	22	21	84	12	31	08	28	29	15	15	0.9	95	95	95
SakSTAR(V112A V113A)	64	130	42	16	39	10	12	34	58	13	80	0.3	15	43	23	30	0.8	95	95	95
SakSTAR(D115A S117A)	80	54	13	14	41	15	15	17	34	19	07	<0.1	15	48	26	13	0.9	95	95	95
SakSTAR(D115A E118A H119A)		32	125	32	34	21	87	13	99	23	93	12	10	24	21	90	1.8	94	95	95
SakSTAR(L116A S117A)	25	<5	44	35	36	33	42	160	29	220	<0.1	<0.1	0.5	41	49	35	1.6			
SakSTAR(H119A K121A)		130	180	24	11	26	29	25	14	29	12	0.52	12	11	29	20	1.2			
SakSTAR(I120A)	26	75	23	26	51	17	16	30	98	25	90	69	30	15	51	52	1.0	93	95	95
SakSTAR(N122A)	5	19	NT															95	93	95
SakSTAR(F123A)	3	<10	28	18	47	11	18	17	32	60	19	<0.1	0.3	53	21	09	1.6	93	90	95
SakSTAR(N126V)	11	51	76	13	20	12	13	10	58	290	98	25	18	80	42	65	0.7	95	95	95
SakSTAR(L127A)	11	54	89	67	18	50	66	25	49	14	84	15	09	19	09	25	1.8	93	94	95
SakSTAR(I128A)	10	20	16	25	48	15	14	38	26	43	82	29	25	20	42	67	0.9	95	93	95
SakSTAR(T129A)	44	190	53	15	23	14	24	21	13	15	42	23	07	10	33	13	1.0	95	95	95
SakSTAR(K130A)	130	280	51	12	32	64	35	22	67	11	15	27	<0.1	<0.1	41	09	0.6	92	74	71
SakSTAR(V131A)	130	70	65	17	29	11	13	19	14	19	29	34	19	13	53	86	0.9	95	95	95
SakSTAR(V132A)	100	130	42	15	26	92	11	33	12	30	19	21	21	36	<0.1	26	0.4	95	95	95
SakSTAR(I133A)	1	99	94	15	19	78	78	24	60	91	86	14	056	64	16	16	0.9	95	95	95
SakSTAR(E134A K135A K136A)		74	122	21	67	25	25	>18	>25	>15	>12	17	02	11	094	60	261			
SakSTAR(K135A)	54	410	52	12	11	79	11	20	11	11	38	20	16	69	37	19	0.9	95	95	95
SakSTAR(K136A)		180	76	18	56	12	13	54	53	12	16	32	35	86	39	18	0.5	91	83	95

LE: expression level below 3 mg/L

**Table 4: Mutagenesis of S34, G36 and H43: Association constants ( $K_A \times 10^7 \text{mol/L}^{-1}$ ) for binding to insolubilized murine monoclonal antibodies (Mab) and absorption (percent) of antibodies of immunized patient plasma**

Variant		Exp (mg/L)	Spec Act (kU/mg)	murine MAbs										SakSTAR patient plasma									
				Epitope cluster I			Epitope cluster II				Epitope cluster III			Pool	Subpool B	Subpool C							
				7G11	26A2	30A2	2B12	3G10	18F12	14H5	28H4	32B2	7F10				7H11	25E1	40C8	24C4	1A10		
SakSTAR				120	9.3	1.1	2.9	7.8	11	18	7.4	19	7.7	2.4	4.0	14	5.4	2.9	0.6	95	95	95	
SakSTAR(S34G.G36R.H43R)				120	10	1.4	3.3	7.5	11	<0.1	<0.1	<0.1	20	2.7	<0.1	<0.1	<0.1	0.15	1.7	87	76	75	
SakSTAR(S34A)				29	110	17	2.4	4.6	9.5	11	28	11	22	15	2.9	3.1	8.8	2.0	0.2	95	95	93	
SakSTAR(G36A)				14	72	3.5	9.8	1.5	5.7	6.5	5.2	4.2	17	9.2	1.4	<0.1	<0.1	0.9	5.0	1.0	86	83	78
SakSTAR(G36E)				12	66	3.1	8.7	1.4	4.8	4.7	12	2.8	6.1	7.6	1.0	<0.1	<0.1	<0.1	3.4	1.1	89	83	72
SakSTAR(G36K)				14	88	9.9	2.3	3.1	8.3	9.8	21	3.9	1.3	15	3.0	<0.1	<0.1	<0.1	2.6	1.2	88	80	69
SakSTAR(G36L)				15	92	3.6	11	1.8	6.1	6.1	16	1.4	6.4	12	1.3	<0.1	<0.1	<0.1	0.6	1.1	93	85	72
SakSTAR(G36N)				10	91	8.7	10	1.6	6.1	6.2	33	3.3	7.8	7.9	1.8	<0.1	<0.1	1.0	0.3	0.5	86	80	75
SakSTAR(G36Q)				21	92	10	12	1.8	6.7	6.5	23	3.8	7.5	7.3	1.5	<0.1	<0.1	<0.1	0.1	0.4	87	84	73
SakSTAR(G36R)				45	100	11	2.4	3.3	10	10	27	4.6	14	20	3.4	<0.1	<0.1	<0.1	3.1	1.2	89	81	70
SakSTAR(H43A)				17	69	15	28	9.7	18	7.6	<0.1	<0.1	<0.1	9.1	1.5	2.0	23	7.8	7.2	1.6	95	95	95
SakSTAR(H43R)				43	120	15	11	2.7	7.6	11	<0.1	<0.1	<0.1	13	6.4	0.7	18	6.7	5.7	1.4	95	95	95
SakSTAR(S34G.G36R)				45	90	3.1	12	2.3	4.8	4.2	13	8.3	2.4	9.1	1.9	<0.1	<0.1	<0.1	<0.1	0.6	92	83	69
SakSTAR(S34G.G36R.H43R.K74A)				1	12	12	4.7	3.8	7.4	9.4	<0.1	0.1	0.6	2.5	2.3	<0.1	<0.1	0.5	0.8	1.7	67	56	83
SakSTAR(S34G.G36R.K74A)				15	26	4.0	2.1	<0.1	1.8	0.6	10	2.2	1.5	1.3	1.8	<0.1	<0.1	<0.1	0.3	2.2	59	28	68
SakSTAR(K35G.G36R.H43D)				32	6	1.8	1.5	1.6	2.0	8.9	<0.1	<0.1	<0.1	7.1	1.3	<0.1	<0.1	<0.1	<0.1	0.9	82	75	72
SakSTAR(G36R.K74A)				40	35	19	7.0	0.2	4.3	2.0	53	2.7	2.8	1.9	4.4	<0.1	<0.1	<0.1	1.2	1.0	48	33	58
SakSTAR(G36R.K74R)				68	150	4.7	1.7	1.8	1.1	8.0	1.6	6.0	6.4	3.0	1.6	<0.1	<0.1	<0.1	0.2	0.8	81	54	73
SakSTAR(G36R.K74A.N95A)				11	25	6.1	2.9	2.9	2.4	<0.1	11	5.7	1.2	5.3	4.1	<0.1	<0.1	<0.1	<0.1	0.9	53	32	63
SakSTAR(G36R.K74A.K135R)				20	33	5.8	3.4	<0.1	1.7	0.7	26	1.6	1.4	1.7	1.2	<0.1	<0.1	<0.1	0.4	0.5	64	32	68
SakSTAR(G36R.K74R.K135R)				48	75	6.1	1.7	5.8	1.0	3.3	31	1.4	1.3	5.7	2.3	<0.1	<0.1	<0.1	0.3	0.8	77	49	68

**Table 5: Mutagenesis of K35, Y73, K74, E80/D82, N95, K130, V132 and K135: Association constants ( $K_A \times 10^7 \text{ mol/L}^{-1}$ ) for binding to insolubilized murine monoclonal antibodies (Mab) and absorption (percent) of antibodies of immunized patient plasma**

murine monoclonal antibodies (Table 1)																				
V <sub>max</sub> unit	Exp	Spec Act (kU/mg)	Epitope cluster I					Epitope cluster II					Epitope cluster III					SakSTAR patient plasma		
			17G11	26A2	30A2	2B12	3G10	18F12	14H5	28H4	32B2	7F10	7H11	25E1	40C8	24C4	1A10	Pool	Subpool B	Subpool C
SakSTAR		120	9.1	1.3	2.9	7.8	11	18	7.4	19	7.7	2.4	4.0	1.4	5.4	2.9	0.6	95	95	95
SakSTAR(S34G,G36R,H43R)		120	10	1.4	3.1	7.5	11	<0.1	<0.1	<0.1	20	2.7	<0.1	<0.1	<0.1	0.15	1.7	87	76	75
SakSTAR(K35A)	230	6.4	1.4	3.1	8.0	7.4	31	11	12	11	2.6	<0.1	1.3	0.2	1.7	0.8	91	88	95	
SakSTAR(K35E)	75	4.8	10	0.6	2.7	2.7	21	5.7	9.1	4.8	1.0	<0.1	<0.1	<0.1	1.5	0.4	95	95	92	
SakSTAR(K35Q)	9	3.2	9.5	1.3	5.2	5.4	22	2.7	9.3	8.5	1.2	0.5	1.7	<0.1	1.9	1.0	95	95	95	
SakSTAR(V73A)	20	<5	9.5	<0.1	<0.1	4.8	33	7.3	11	8.9	2.0	0.5	9.0	2.5	1.4	0.8	63	44	93	
SakSTAR(V71F)	6	11	7.9	1.2	1.3	2.1	8.0	24	15	19	1.5	2.6	1.6	6.2	3.5	7.2	1.4	93	95	95
SakSTAR(V73H)	30	<5	7.3	1.5	<0.1	<0.1	1.4	20	7.5	1.7	4.2	4.3	3.5	6.8	5.9	9.7	1.3	76	65	95
SakSTAR(V71L)	33	<5	11	<0.1	<0.1	<0.1	84	19	26	5.3	3.4	4.2	7.1	4.2	3.4	1.0	81	60	94	
SakSTAR(V73S)	11	<5	7.0	1.5	0.59	0.6	3.0	21	8.3	1.4	21	3.2	2.2	6.7	1.3	1.3	0.7	86	69	95
SakSTAR(V71W)	6	27	4.8	9.0	4.6	4.6	11	8.4	4.5	11	8.0	2.7	2.9	5.0	3.0	3.8	1.3	73	53	93
SakSTAR(K74A)	80	69	4.4	2.7	0.2	2.2	1.1	17	5.2	1.4	7.6	2.2	2.0	6.8	3.3	1.8	0.9	64	58	95
SakSTAR(K74E)	12	<5	2.2	0.6	<0.1	0.7	0.1	1.4	1.8	7.5	9.3	1.2	2.0	3.0	1.2	0.6	1.0	83	43	90
SakSTAR(K74N)	9	39	2.9	4.7	1.1	3.3	1.7	10	1.6	4.8	1.3	1.5	1.9	4.0	1.8	1.4	0.9	63	46	95
SakSTAR(K74Q)	64	110	5.3	5.8	<0.1	2.5	1.1	24	5.9	1.2	5.4	1.1	2.0	6.2	2.3	2.0	0.4	70	62	94
SakSTAR(K74R)	44	150	2.1	7.5	2.0	4.1	4.2	24	6.9	8.0	8.3	1.3	2.2	7.8	3.2	2.1	0.5	75	70	95
SakSTAR(E80A,D82A)	130	7.3	1.2	2.1	6.5	5.9	7.9	6.1	8.4	7.8	1.9	<0.1	<0.1	<0.1	<0.1	0.4	89	83	92	
SakSTAR(E80A)	160	1.3	1.3	3.3	7.9	10	35	7.4	1.7	8.6	2.1	<0.1	<0.1	1.6	3.6	<0.1	1.7	94	93	95
SakSTAR(D82A)	160	1.7	1.2	4.8	7.3	11	31	7.8	1.7	1.2	2.7	<0.1	<0.1	0.2	<0.1	<0.1	2.3	95	93	95
SakSTAR(N95A)	25	260	10	1.8	4.0	10	11	50	1.3	1.4	4.9	2.3	1.7	7.3	4.7	2.9	0.8	95	94	95
SakSTAR(N95E)	12	79	2.8	8.5	1.3	5.2	5.4	17	2.7	10	5.2	1.1	0.5	4.0	1.7	1.8	0.6	95	92	95
SakSTAR(N95G)	20	160	4.1	11	1.5	6.8	7.6	36	3.3	1.5	3.6	1.5	0.7	5.8	2.7	2.7	0.9	95	90	95
SakSTAR(N95K)	54	180	9.5	1.4	3.2	9.0	11	51	5.0	1.8	4.8	2.5	1.6	8.3	2.9	4.8	1.1	95	95	95
SakSTAR(N95R)	LE																			

Table 5 - cont'd: Mutagenesis of K35, Y73, K74, E80/D82, N95, K130, V132 and K135: Association constants ( $K_A \times 10^7 \text{ mol/L}^{-1}$ ) for binding to insolubilized murine monoclonal antibodies (Mab) and absorption (percent) of antibodies of immunized patient plasma

V <sub>ARI</sub> ant	Exp (mg/L)	Spec Act (kU/mg)	murine MAbs										SAKSTAR patient plasma							
			Epitope cluster I					Epitope cluster II					Epitope cluster III					Subpool B		Subpool C
			7G11	26A2	30A2	28B2	3G10	18F12	14H5	28H4	32B2	7F10	7H11	25E1	40C8	24C4	1A10	Pool		
SAKSTAR(K130A)		280	5.1	1.2	3.2	6.4	3.5	22	6.7	11	15	2.7	<0.1	<0.1	4.1	0.9	0.6	92	74	71
SAKSTAR(K130T)		290	7.8	1.4	3.1	8.0	5.6	43	5.4	9.7	17	2.7	<0.1	<0.1	3.4	0.5	0.5	92	84	82
SAKSTAR(V132A)	102	130	4.2	1.5	2.6	9.2	11	33	12	30	19	2.1	2.1	3.6	<0.1	2.6	0.4	95	95	95
SAKSTAR(V132L)	126	120	4.8	1.4	2.3	8.0	9.7	67	10	48	20	2.5	2.0	1.1	<0.1	4.8	0.4	95	95	95
SAKSTAR(V132T)	78	140	4.5	1.3	2.4	7.8	9.0	39	10	25	16	2.0	1.1	2.0	<0.1	<0.1	0.4	95	95	95
SAKSTAR(V132N)	16	150	4.5	1.1	1.7	7.0	7.2	21	17	20	15	2.0	1.5	1.3	<0.1	4.2	0.7	95	95	95
SAKSTAR(V132R)	76	230	5.4	1.2	0.8	3.3	3.2	23	5.5	7.8	4.8	1.1	2.1	3.4	<0.1	0.4	0.4	95	95	95
SAKSTAR(K135A)	54	410	5.2	1.2	1.1	7.9	11	20	11	11	3.8	2.0	1.6	6.9	3.7	1.9	0.9	95	95	95
SAKSTAR(K135F)	68	64	3.9	6.3	1.0	6.1	4.1	13	4.5	8.4	16	2.1	1.6	5.8	1.9	1.5	0.9	95	95	95
SAKSTAR(K135R)	54	230	4.0	1.4	1.4	9.3	5.0	11	8.1	1.3	2.3	1.9	2.4	2.4	2.6	2.1	0.5	95	95	95
SAKSTAR(K15A, K74A)	20	130	NT															84	50	95
SAKSTAR(Y73A, K74A)	24	<5	18	<0.1	<0.1	<0.1	<0.1	19	6.7	23	9.9	3.2	2.7	1.3	4.0	1.6	1.1	47	28	87
SAKSTAR(Y73F, K74A)	21	6	17	6.7	<0.1	2.1	<0.1	42	19	21	20	4.9	1.9	2.8	6.6	2.1	0.8	51	34	90
SAKSTAR(I60A, K74A, N95A)		84	5.3	2.7	<0.1	2.5	5.3	17	7.2	9.6	3.4	2.3	2.1	6.2	2.7	2.9	0.8	60	47	95
SAKSTAR(N95A, K135R)	120	240	4.9	1.3	1.9	9.0	9.9	18	12	15	2.5	1.5	1.8	5.9	3.6	3.7	0.8	95	95	95
SAKSTAR(K130T, K135R)	15	280	1.7	1.0	1.6	7.2	8.0	13	7.7	4.5	3.7	1.6	<0.1	<0.1	2.4	0.4	0.6	89	60	73

Table 6: Combination mutants of SakSTAR(K130T,K135R) with K35A, G36R, E65X,K74X and selected other amino acids

N. strain		Exp (mg/mL)	Spec Act (kU/mg)	Epitope cluster I					Epitope cluster II					Epitope cluster III					SakSTAR patient plasma				Code		
				Epitope cluster I					Epitope cluster II					Epitope cluster III					SakSTAR patient plasma						
				17G1T	36A2	10A2	2812	3G10	18F12	14H5	28H4	37B2	7F10	7H11	23E1	40C8	23C4	1A10	Pool 10	Subpool B	Subpool C	Pool 40			
				15	280	3.7	10	16	7.2	8.0	13	7.7	4.3	3.7	1.6	<0.1	<0.1	2.4	0.4	0.6	89	60	73	90	SY2
				26	220	7.2	18	2.8	11	14	17	3.9	8.9	5.1	1.6	<0.1	<0.1	<0.1	<0.1	1.1	79	65	69	-	SY3
				18	310	7.3	27	5.9	16	18	17	3.7	11	5.1	1.5	<0.1	<0.1	3.2	0.7	0.9	76	49	69	78	SY4
				64	190	4.0	7.2	3.0	7.7	0.9	30	5.7	7.2	8.4	1.8	<0.1	<0.1	2.7	0.7	1.0	50	25	67	62	SY41
				5	210	7.6	26	5.7	17	19	17	4.7	10	5.0	1.4	<0.1	<0.1	<0.1	<0.1	0.8	73	44	69	75	SY5
				88	120	5.5	7.3	0.8	11	6.4	15	11	7.1	6.1	2.6	<0.1	<0.1	<0.1	<0.1	0.8	51	25	63	54	SY42
				29	160	5.6	8.8	2.2	10	10	<0.1	<0.1	<0.1	6.3	2.0	<0.1	<0.1	<0.1	<0.1	0.9	72	39	69	-	SY9
				40	76	4.8	5.9	0.5	11	1.5	18	8.7	7.5	5.1	2.1	<0.1	<0.1	<0.1	<0.1	1.1	52	25	65	61	SY43
				46	170	11	9.0	2.1	11	11	19	66	12	5.7	2.3	<0.1	<0.1	3.5	1.2	0.6	45	16	77	55	SY48
				80	83	4.7	12	1.2	16	21	27	12	10	6.9	2.6	<0.1	<0.1	<0.1	<0.1	0.7	44	18	65	46	SY44
				17	71	5.7	10	1.8	11	1.2	21	8.4	6.9	4.6	1.8	<0.1	<0.1	<0.1	<0.1	1.0	41	14	64	50	SY59
				60	96	5.6	6.0	2.0	3.6	4.9	21	8.6	8.8	2.8	2.5	<0.1	<0.1	3.5	1.7	1.1	51	13	66	56	SY51
				40	150	6.7	18	2.1	15	16	9.0	3.1	4.1	6.3	2.3	<0.1	<0.1	3.8	0.9	0.6	53	29	67	63	SY49
				54	130	2.4	4.9	<0.1	7.4	3.8	19	8.7	7.6	4.7	1.9	<0.1	<0.1	3.5	1.3	1.5	56	32	69	61	SY55
				12	210	6.2	13	1.8	10	11	11	3.9	5.0	6.8	2.3	<0.1	<0.1	3.1	1.1	0.8	49	21	64	59	SY63
				36	46	7.7	<0.1	<0.1	<0.1	<0.1	10	3.9	3.2	7.4	2.6	<0.1	<0.1	4.6	1.3	0.6	43	15	62	55	SY66
				35	67	7.0	<0.1	<0.1	<0.1	<0.1	13	5.4	4.9	6.6	2.5	<0.1	<0.1	3.2	1.3	1.0	49	29	63	57	SY67
				19	78	4.3	24	<0.1	2.7	5.6	20	9.6	7.5	5.6	2.2	<0.1	<0.1	1.7	1.1	1.2	37	12	57	50	SY68
				28	240	5.6	5.4	0.5	7.5	5.1	21	8.4	14	6.1	2.4	<0.1	<0.1	4.1	2.6	0.7	57	27	78	65	SY56
				60	230	6.0	14	2.4	1.5	17	9.0	3.3	5.8	6.3	2.3	<0.1	<0.1	4.3	2.0	0.6	51	32	73	58	SY69
				46	300	4.1	4.4	0.8	6.6	3.8	18	8.5	7.8	5.2	2.1	<0.1	<0.1	2.4	0.7	0.9	55	29	64	59	SY57
				88	170	5.3	8.9	1.7	7.7	11	16	3.4	5.8	6.2	2.4	<0.1	<0.1	3.5	2.4	0.7	55	27	79	55	SY70
				68	170	4.5	4.9	0.4	6.0	5.2	9.0	4.9	4.3	5.6	2.4	<0.1	<0.1	<0.1	<0.1	0.6	51	20	63	56	SY58
				16	170	6.2	13	1.8	12	14	7.3	2.5	3.8	<0.1	<0.1	<0.1	<0.1	4.1	1.9	0.5	51	27	69	57	SY71

**Table 6 - cont'd: Combination mutants of SakSTAR(K130T,K135R) with K35A, G36R, E65X,K74X and selected other amino acids**

Table 6 - Cont'd. Combination mutants																					
V <sub>mutant</sub>	Exp mg/L	Spec. Act (kU/mg)	mutant NAbx										Epitope cluster III				SakSTAR patient plasma				Code
			Epitope cluster I					Epitope cluster II					Epitope cluster III				Pool 10		Subpool C		
			7G11	26A2	30A2	2812	3G10	18F12	14H5	28R4	32B2	7F10	7H11	23E1	40C8	24C4	1A10	Pool 10	Subpool B	Subpool C	Pool 40
	40	220	61	14	19	17	15	90	33	58	63	23	<0.1	<0.1	4.1	2.2	0.5	52	29	74	58

Association constants  $\geq 10$ -fold lower and antibody absorption  $\leq 60$  percent of wild-type SakSTAR are represented in bold type;  $\geq 100,000$  HU/mg represented in bold type. NT: not tested.

**Table 7: Combination mutants of SakSTAR(E80A,D82A,K130T,K135R) with K35A, G36R, E65X, K74X, and selected other amino acids**

Exp (mg/mL)		Spec. Act (kU/mg)	mouse MAbs										Epitope cluster III			SakSTAR patient plasma			Code	
			Epitope cluster I					Epitope cluster II					Epitope cluster III			SakSTAR patient plasma				
			17G11	26A2	30A2	2812	3G10	18F12	14H5	28H4	32B2	7F10	7H11	23E1	40C8	24C4	1A1	Pool 10		Subpool B
20	250	8.4	17	3.1	11	13	18	4.3	11	5.5	1.5	<0.1	<0.1	<0.1	<0.1	0	80	64	68	SY6
4	220	5.3	31	2.8	18	11	89	8.4	21	5.9	1.8	<0.1	<0.1	<0.1	<0.1	0.6	74	34	69	SY7
27	110	5.1	6.5	1.2	7.9	7.3	28	6.4	19	4.6	1.7	<0.1	<0.1	<0.1	<0.1	0.8	46	17	60	SY15
70	160	5.9	5.7	6.4	2.7	18	19	16	8.2	9.0	1.8	<0.1	<0.1	<0.1	<0.1	0.8	66	34	66	SY17
22	140	5.4	50	2.9	49	21	31	11	3.7	3.4	1.8	<0.1	<0.1	<0.1	<0.1	0.6	43	11	68	SY19
1	110	1.2	1.2	4.6	4.4	1.5	19	5.0	1.5	6.6	1.4	<0.1	<0.1	<0.1	<0.1	0.4	35	12	60	SY20
10	94	7.2	9.3	9.9	5.6	3.2	22	15	8.2	11	2.0	<0.1	<0.1	<0.1	<0.1	1.0	58	24	69	SY21
250	250	5.6	54	2.9	38	2.5	36	6.5	18	5.5	1.9	<0.1	<0.1	<0.1	<0.1	0.9	75	33	68	SY10
40	140	8.2	10	1.3	5.7	5.5	24	22	11	1.5	2.2	<0.1	<0.1	<0.1	<0.1	1.0	51	17	66	SY18
88	120	8.5	12	1.1	7.0	3.6	43	13	12	7.9	2.4	<0.1	<0.1	<0.1	<0.1	0.9	60	29	67	SY21
55	140	9.0	16	1.6	9.9	5.9	18	4.5	4.3	10	2.5	<0.1	<0.1	<0.1	<0.1	0.8	54	22	63	SY22
24	110	2.4	17	2.9	1.3	1.2	16	7.8	16	4.5	1.6	<0.1	<0.1	<0.1	<0.1	0.7	75	64	62	SY13
92	62	8.2	2.3	3.8	8.4	1.9	30	11	1.3	3.6	3.1	<0.1	<0.1	<0.1	<0.1	1.2	51	13	66	SY53
84	110	7.0	3.9	2.6	4.1	7.2	29	28	14	6.8	2.2	<0.1	<0.1	<0.1	<0.1	0.9	43	13	64	SY30
54	120	5.1	16	3.2	1.4	3.7	16	20	2.4	5.7	1.7	<0.1	<0.1	<0.1	<0.1	1.0	43	21	64	SY47
56	140	5.1	6.8	2.6	9.5	1.7	28	17	14	8.4	2.1	<0.1	<0.1	<0.1	<0.1	1.0	35	8	58	SY46



**Table 7 - cont'd: Combination mutants of SakSTAR(E80A,D82A,K130T,K135R) with K35A, G36R, E65X, K74X, and selected other amino acids**

Variant	Exp (mg/mL)	Spec Act (kU/mg)	Epitope cluster I					murine MAbs Epitope cluster II					Epitope cluster III					SakSTAR patient plasma				Code
			17G11	26A2	30A2	2812	3G10	18H12	14H5	28H4	32B2	7F10	7H11	25E1	40C8	24C4	1A10	Pool 10	Subpool B	Subpool C	Pool 40	
SakSTAR(K74R,E80A,D82A,S103A,K130T,K135R)	32	160	1.9	2.2	3.8	14	4.3	26	6.9	7.4	5.0	1.7	<0.1	<0.1	<0.1	<0.1	0.9	67	32	69	70	SY24
SakSTAR(K15A,E65D,K74R,E80A,D82A,E108A,K109A,K130T,K135R)	9.0	89	5.8	3.0	2.6	26	10	13	16	13	3.2	1.8	<0.1	<0.1	<0.1	<0.1	0.5	55	10	63	47	SY12
SakSTAR(K35A,E65D,K74R,E80A,D82A,E108A,K130T,K135R)	20	91	6.4	2.0	5.0	15	3.9	22	17	7.4	2.4	1.9	<0.1	<0.1	<0.1	<0.1	0.9	44	8	70	53	SY32
SakSTAR(E65D,K74R,E80A,D82A,E108A,K110T,K135R)	4	90	8.1	6.7	6.9	4.3	29	19	31	11	14	2.1	<0.1	<0.1	<0.1	<0.1	1.0	52	11	69	-	SY33
SakSTAR(K35A,E65D,K74R,E80A,D82A,K109A,K130T,K135R)	42	84	5.5	1.8	5.3	14	1.6	18	12	7.7	10	1.7	<0.1	<0.1	<0.1	<0.1	1.0	43	6	61	50	SY36
SakSTAR(E65D,K74R,E80A,D82A,K109A,K130T,K135R)	60	130	9.7	6.6	6.8	4.2	28	11	32	12	17	2.3	<0.1	<0.1	<0.1	<0.1	0.9	56	10	64	53	SY37
SakSTAR(K35A,E65D,K74R,E80A,D82A,K110T,K135R,K136A)	28	81	4.5	1.2	3.3	11	1.7	22	13	7.6	4.9	1.6	<0.1	<0.1	<0.1	<0.1	0.8	40	14	52	40	SY34
SakSTAR(E65D,K74R,E80A,D82A,K130T,K135R,K136A)	60	100	6.8	5.8	4.4	4.5	15	33	32	14	7.9	2.0	<0.1	<0.1	<0.1	<0.1	0.8	46	28	67	45	SY35
SakSTAR(E65Q,K74Q,D82A,S84A,K130T,K135R)		170	NT															45	21	60	45	SY30N
SakSTAR(K15A,E65D,K74R,E80A,D82A,K86A,K130T,K135T)	68	86	4.4	2.0	5.5	15	1.5	15	12	6.4	6.7	1.9	<0.1	<0.1	<0.1	<0.1	1.0	36	7	60	55	SY40
SakSTAR(K35A,K74Q,E80A,D82A,K130T,K135R)	72	120	6.1	3.4	2.5	3.0	5.9	38	14	9.8	6.8	1.9	<0.1	<0.1	<0.1	<0.1	0.8	49	16	64	48	SY28
SakSTAR(K35A,E65D,K74R,E80A,D82A,K130T,K135R)	54	190	8.1	7.5	6.9	5.5	25	37	34	14	7.7	2.3	<0.1	<0.1	<0.1	<0.1	1.0	56	28	68	55	SY29
SakSTAR(K35A,E65D,K74R,E80A,D82A,V132R,K135R)	11	55	6.7	2.3	5.3	17	2.3	47	19	19	5.1	2.0	<0.1	<0.1	<0.1	<0.1	1.1	53	20	88	62	SY61
SakSTAR(K35A,E65D,K74R,E80A,D82A,T129A,K135R)	11	61	7.0	1.3	5.1	31	12	27	12	11	6.7	2.5	<0.1	<0.1	<0.1	<0.1	1.9	56	18	79	60	SY62
SakSTAR(K35A,E65D,K74R,E80A,D82A,T129A,K135A)	23	21	6.9	2.7	5.8	32	20	29	6.6	9.7	5.4	2.1	<0.1	<0.1	<0.1	<0.1	0.9	56	17	91	60	SY64

Association constants  $\geq$  10-fold lower and antibody absorption  $\leq$ 60 percent of wild-type SakSTAR are represented in bold type;  $\geq$  100,000 HU/mg represented in bold type. NT: not tested.

**Table 8: SakSTAR variants with intact specific activity ( $\geq 100$  kHU/mg) and  $\leq 50$  percent absorption of human antibodies elicited by treatment with wild-type SakSTAR**

Variant	Spec. Act. (kU/mg)	SakSTAR patient plasma			Code
		Pool 10	Subpool B	Subpool C	
SakSTAR(K74Q,K130T,K135R)	190	50	25	67	62
SakSTAR(E65A,K74Q,K130T,K135R)	170	45	16	77	55
SakSTAR(E65Q,T71S,K74Q,K130T,K135R)	210	49	21	64	59
SakSTAR(E65Q,K74Q,E118A,K130A,K135R)	180	50	28	72	58
SakSTAR(E65Q,K74Q,N95A,E118A,K130A,K135R)	190	48	27	74	58
SakSTAR(K35A,E65Q,K74Q,K130A,K135R)	110	49	26	63	45
SakSTAR(E65Q,K74Q,K109A,K130T,K135R)	210	50	22	68	51
SakSTAR(K74Q,E80A,D82A,K130T,K135R)	110	46	17	60	48
SakSTAR(E65D,K74R,E80A,D82A,K130T,K135R)	140	43	11	68	57
SakSTAR(E65S,K74R,E80A,D82A,K130T,K135R)	110	35	12	60	-
SakSTAR(E65D,K74R,E80A,D82A,K130T,K135R,K136A)	100	46	28	67	45
SakSTAR(K35A,K74Q,E80A,D82A,K130T,K135R)	120	49	16	64	48
SakSTAR(E65D,K74Q,E80A,D82A,K130T,K135R)	110	43	13	64	42
SakSTAR(E65Q,K74Q,E80A,D82A,K130T,K135R)	120	43	21	64	42
SakSTAR(E65Q,K74Q,D82A,S84A,K130T,K135R)	170	45	21	60	45
SakSTAR(K35A,E65D,K74Q,E80A,D82A,K130T,K135R)	140	35	8	58	40
SakSTAR(T21A,K35A,E65Q,K74Q,K130A,K135R)	110	50	26	72	50
SakSTAR(E65Q,K74Q,K109A,K121A,K130A,K135R)	140	50	31	73	52
SakSTAR(E65Q,K74Q,D82A,S84A,K109A,K130A,K135R)	180	43	20	62	44

**Table 8 - cont'd: SakSTAR variants with intact specific activity ( $\geq 100$  kHU/mg) and  $\leq 50$  percent absorption of human antibodies elicited by treatment with wild-type SakSTAR**

Variant	Spec. Act. (kU/mg)	SakSTAR patient plasma				Code
		Pool 10	Subpool B	Subpool C	Pool 40	
SakSTAR(E65Q,K74Q,N95A,E118A,K130A,K135R,V137A)	120	45	30	74	60	SY93
SakSTAR(E65Q,K74Q,N95A,E118A,K130A,K135R,K136A,V137K)	1,400	37	16	70	54	SY94
SakSTAR(E65Q,K74Q,D82A,S84A,E108A,K109A,K130A,K135R)	110	46	26	63	41	SY95

Antibody absorption  $\leq 60$  percent of wild-type SakSTAR are represented in bold type;  $\geq 100,000$  HU/mg represented in bold type.

**Table 2: Fibrinolytic properties of selected SakSTAR variants in human plasma in vitro**

Compound	Fibrinolytic potency (C50 in µg/mL)	Residual fibrinogen at C50 (% of baseline)	Fibrinogenolytic potency (C50 in µg/mL)	Code
SakSTAR	0.18 ± 0.01	93 ± 3.5	24 ± 3.6	SY15
SakSTAR(K74Q,E80A,D82A,K130T,K135R)	0.15 ± 0.01	97 ± 3.0	14 ± 3.2	SY19
SakSTAR(E65D,K74R,E80A,D82A,K130T,K135R)	0.24 ± 0.04	94 ± 10	29 ± 3.1	SY46
SakSTAR(K35A,E65D,K74Q,E80A,D82A,K130T,K135R)	0.11 ± 0.01	92 ± 3.0	20 ± 2.0	SY93
SakSTAR(E65Q,K74Q,N95A,E118A,K130A,K135R,K136A,∇137K)	0.13	91		

The data represent mean ± SD of 3 experiments.  
C<sub>50</sub>: amount of wild type or variant SakSTAR required for 50% clot lysis or 50% fibrinogen breakdown in 2 hrs.

**Table 10: Pharmacokinetic parameters of the disposition of staphylokinase-related antigen from plasma following bolus injection of SakSTAR variants (100 µg/kg) in hamsters.**

Variant	C <sub>0</sub> (µg/mL)	A (µg/mL)	B (µg/mL)	t <sub>1/2</sub> (α) (min)	t <sub>1/2</sub> (β) (min)	VC (mL)	AUC (µg.min.mL <sup>-1</sup> )	Cl <sub>p</sub> (mL.min <sup>-1</sup> )
SakSTAR	0.8 ± 0.1	0.6 ± 0.1	0.2 ± 0.0	2.8	7.0	13 ± 1.0	4.6 ± 0.4	2.2 ± 0.2
SakSTAR(K74Q,E80A,D82A,K130T,K135R)	0.5 ± 0.1	0.4 ± 0.1	0.1 ± 0.0	2.0	10	20 ± 2.2	2.5 ± 0.3	4.1 ± 0.5
SakSTAR(E65D,K74R,E80A,D82A,K130T,K135R)	0.6 ± 0.0	0.5 ± 0.0	0.1 ± 0.0	2.0	10	16 ± 1.1	2.8 ± 0.2	3.7 ± 0.3
SakSTAR(K35A,E65DK74Q,E80A,D82A,K130T,K135R)	1.1 ± 0.1	1.0 ± 0.1	0.1 ± 0.0	2.0	24	9.6 ± 0.7	6.4 ± 0.5	1.6 ± 0.1

Data are mean ± SEM of 4 experiments.

**Table 11: Baseline characteristics and treatment outcome of the patients with peripheral arterial occlusion treated with SakSTAR, SakSTAR(K74Q,E80A,D82A,K130T,K135R) or SakSTAR(E65D,K74R,E80A,D82A,K130T,K135R)**

Compound Patient Id.	Gender	Age (yrs)	Clinical ischemia	Locus of occlusion	Age of occlusion (days)	Length of occlusion (cm)	Recanalization by thrombolysis	Total dose of thrombolytic agent (mg)	Total duration of infusion (hrs)	Additional therapy
<b>SakSTAR</b>										
PUT	M	66	Subacute	Femoro-femoral graft	6	5	Complete	2	23	Stenting left IF artery
VERM	M	73	Acute	Right PA	2	6	Partial	13	23	Right upper leg amputation
GEIV	V	63	Respain	Left SFA	10	5	Complete	8	7	PTA
POL	M	46	Subacute	Right SFA	30	50	Partial	22	29	Lumbar sympathectomy
BUE	F	53	Claudication	Right AF graft	1	15	Complete	10	13	Desobstruction
VUJ	F	75	Subacute	Left FT graft	2	34	Complete	7	10	PTA
REN	M	48	Respain	Right IF graft	4	20	Complete	6.5	5	Left AF graft
COR	V	78	Acute	Left AFS	14	9	Complete	4	3	PTA
MAN	M	67	Respain	Left tibial artery	1	6	Partial	6	5	-
STRA	M	66	Claudication	Right FP graft	14	16	Complete	19	26	-
VANH	M	38	Acute	Left radial artery	4	1	Complete	6	5	-
VANW	F	57	Acute	Right FP graft	1	25	Complete	20	24	New right FP graft
BRA	M	57	Acute	Left FT graft	1	30	Complete	25	43	-
DON	M	60	Claudication	Left FT graft	1	20	Complete	13	19	PTA + stenting
CAM	M	77	Respain	Right SFA graft	8	30	Complete	27	44	FF graft
Mean ± SEM		62 ± 3.1			6.6 ± 2.1	18 ± 3.5		13 ± 2.1	19 ± 3.5	
<b>SakSTAR(K74Q,E80A,D82A,K130T,K135R)</b>										
IMB	M	66	Claudication	Left SFA	30	5	Complete	24	24	PTA
AZY	M	44	Subacute	Right C.I.A.	7	8	Complete	18	23	Stenting
VIN	M	51	Acute	Right E.I.A.	5	70	Complete	24	30	-
STRO	M	53	Claudication	Left FP junction	14	5	Partial	3.5	2	Aspiration thrombectomy, PTA
VERG	M	62	Respain	Left SFA	20	6	Complete	19	25	FP bypass
GIE	M	76	Acute	Right FP bypass	2	15	Complete	8.5	17	-
Mean ± SEM		59 ± 4.7			13 ± 4.3	18 ± 10		16 ± 3.4	20 ± 4.0	
<b>SakSTAR(E65D,K74R,E80A,D82A,K130T,K135R)</b>										
URB	M	57	Subacute	Right E.I.A.	4	8	Complete	8	6	Pseudo aneurysm, right AF graft revision
COM	M	59	Respain	Right AF graft	7	65	Complete	16	22	-
HIAC	M	70	Respain	Left anterior tibial artery	7	15	Complete	12	14	-
DEW	F	76	Respain	SFA	21	6	Complete	6	4	-
VAI	F	65	Subacute	Left PA	25	10	Partial	8	6	Aspiration thrombectomy
FIL	M	76	Claudication	Right SFA	28	8	Complete	24	31	PTA
Mean ± SEM		67 ± 3.4			15 ± 4.3	19 ± 9.4		12 ± 2.8	14 ± 4.4	

AF: aortofemoral; CABG: coronary artery bypass graft; CAD: coronary artery disease; CIA: common iliac artery; COPD: chronic obstructive pulmonary disease; DM: diabetes mellitus; EIA: external iliac artery; FF: femorofibular; FP: femoropopliteal; FT: femorotibial; IA: iliac artery; IF: iliofemoral; occl: occlusion; PA: popliteal artery; PTA: percutaneous transluminal angioplasty; SFA: superficial femoral artery; TA: tibial artery; TF: tibiofibular; SC: subclavian

**Table 12: Absorption with SakSTAR variants of antibodies elicited with SakSTAR variants in patients with peripheral arterial occlusion**

Treatment	Absorbant	Insolubilized compound		
		SakSTAR	SakSTAR(K74Q,E80A,D82A,K130T,K135R)	SakSTAR(E65D,K74R,E80A,D82A,K130T,K135R)
SakSTAR (Pool 40)	SakSTAR	95		
	SakSTAR(K74Q,E80A,D82A,K130T,K135R)	48		
	SalSTAR(E65D,K74R,E80A,D82A,K130T,K135R)	57		
SakSTAR(K74Q,E80A,D82A,K130T,K135R) (Imb., Vin., Ver., Cie.)	SakSTAR	94	95	95
	SakSTAR(K74Q,E80A,D82A,K130T,K135R)	91	93	89
	SalSTAR(E65D,K74R,E80A,D82A,K130T,K135R)	92	94	94
SakSTAR(E65D,K74R,E80A,D82A,K130T,K135R) (Urb.)	SakSTAR	90	88	85
	SakSTAR(K74Q,E80A,D82A,K130T,K135R)	94	95	94
	SalSTAR(E65D,K74R,E80A,D82A,K130T,K135R)	94	95	94

Data represent median values of the percent absorption with 250 nM absorbant, measured by residual binding to insolubilized compound.

**Table 13: Additive substitution mutagenesis of SakSTAR(E65Q,K74Q,K130T, K135R) with selected other amino acids**

Variant	Spec. Act. (kU/mg)	Antibody absorption (percent)	Code
SakSTAR(E65Q,K74Q,K130T,K135R)	150	65	SY49
SakSTAR(E65Q,K74Q,D82A,S84A,K130T,K135R)	170	45	SY50
SakSTAR(E65Q,K74Q,T90A,E99D,T101S,K130A,K135R)	410	51	SY98
SakSTAR(E65Q,K74Q,E108A,K109A,K130T,K135R)	180	50	SY83
SakSTAR(E65Q,K74Q,D82A,S84A,E108A,K109A,K130T,K135R)	110	41	SY95
SakSTAR(E65Q,K74Q,D82A,S84A,E108A,K109A,K130T,K135R,K136A,V137K)	1,500	30	SY118
SakSTAR(E65Q,K74Q,D82A,S84A,T90A,E99D,T101S,E108A,K109A,K130T,K135R,K136A,V137K)	2,900	28	SY128
SakSTAR(K35A,E65Q,K74Q,D82A,S84A,T90A,E99D,T101S,E108A,K109A,K130T,K135R,K136A,V137K)	3,700	24	SY141
SakSTAR(K35A,E65Q,K74R,D82A,S84A,T90A,E99D,T101S,E108A,K109A,K130T,K135R,K136A,V137K)	5,700	31	SY145

Spec. Act.  $\geq$  100 kU/mg is represented in bold type. Absorption of antibodies (in percent) from pooled immunized patient plasma; values  $\leq$ 60% are represented in bold type.



**Table 14: Fibrinolytic properties of SakSTAR variants in human plasma in vitro**

Compound	Fibrinolytic potency (C50 in µg/ml)	Residual fibrinogen at C50 (% of baseline)	Fibrinogenolytic potency (C50 in µg/ml)	Code
SakSTAR	0.18 ± 0.01	93 ± 3.5	24 ± 3.6	
SakSTAR(E65Q,K74Q,D82A,S84A,E108A,K109A,K130T,K135R,K136A,V137K)	0.15 ± 0.02	90 ± 5.0	14 ± 1.0	SY118
SakSTAR(K35A,E65Q,K74Q,D82A,S84A,T90A,E99D,T101S,E108A,K109A,K130T,K135R,K136A,V137K)	0.17 ± 0.01	87 ± 3.0	7 ± 0.6	SY141
SakSTAR(K35A,E65Q,K74R,D82A,S84A,T90A,E99D,T101S,E108A,K109A,K130T,K135R,K136A,V137K)	0.19 ± 0.01	82 ± 3.0	7 ± 0.9	SY145

The data represent mean ± SD of 3 experiments.

C50: amount of wild type or variant SakSTAR required for 50% clot lysis or 50% fibrinogen breakdown in the absence of fibrin in 2 hrs.

**Table 15:** Characteristics of the patients with peripheral arterial occlusion treated with SakSTAR(E65Q,K74Q,D82A,S84A,E108A,K109A,K130T,K135R, K136A,V137K), SakSTAR(K35A,E65Q,K74Q,D82A,S84A,T90A,E99D,T101S,E108A,K109A,K130T,K135R,K136A,V137K) or SakSTAR(K35A,E65Q, K74R,D82A,S84A,T90A,E99D,T101S, E108A,K109A,K130T,K135R,K136A,V137K)

Compound Patient Id.	Gen-der	Age (yrs)	Clinical ischemia	Risk factors Relevant history	Current Smoking	Locus of occlusion	Age of occlusion (days)	Length of occlusion (cm)
<b>SakSTAR(E65Q,K74Q,D82A,S84A,E108A,K109A,K130T,K135R,K136A,V137K) (SY118)</b>								
VCL	M	69	Acute	Hypertension, ischemic heart disease, ABF graft	-	Left AF graft	10	14
REN	M	76	Subacute	Hypercholesterolemia	-	Right PA	18	14
HOL	M	69	Acute	Hypertension, hypercholesterolemia, right bypass	+	Right FT bypass	16	3.0
PAR	M	79	Pain, swelling	-	-	Left popliteal to common femoral vein	50	8.0
FRA	M	60	Subacute	Ischemic heart disease, left FP graft	+	Left FP graft	30	14
MAC	V	73	Acute	Hypertension, ABF graft	-	Left branch ABF graft	20	10
Mean $\pm$ SEM		71 $\pm$ 2.7					21 $\pm$ 6.9	11 $\pm$ 1.8
<b>SakSTAR(K35A,E65Q,K74Q,D82A,S84A,T90A,E99D,T101S,E108A,K109A,K130T,K135R,K136A,V137K) (SY141)</b>								
VERH	F	52	Claudication	Hypertension, hypercholesterolemia, right IF endoprosthesis	+	Right IA	14	12
DUB	M	54	Claudication	Hypertension, stenosing left, right IA	+	Right EIA	30	18
VAP	M	46	Claudication	Hypertension, hypercholesterolemia, stenosing left + right IA	-	Aortabifurcation	21	25
WYN	M	43	Claudication	CAD; hypercholesterolemia; stenosing left FP graft	+	Left FP graft	5.0	30
HOR	M	57	Acute	Hypertension; left FP graft	+	Left CIA, left FP graft	7.0	>60
AND	M	75	Acute	Diabetes; hypertension; cardiac valve replacements	-	Left SF artery	3.0	10
Mean $\pm$ SEM		55 $\pm$ 4.6					13 $\pm$ 4.3	19 $\pm$ 3.5
<b>SakSTAR(K35A,E65Q,K74R,D82A,S84A,T90A,E99D,T101S,E108A,K109A,K130T,K135R,K136A,V137K) (SY145)</b>								
LIN	F	48	Rest pain	Hypertension, ischemic heart disease	+	Right SF artery	7.0	16
DEL	M	68	Claudication	Hypertension	+	Left PA	21	6.0
LAM	M	71	Acute	FP graft	-	Right FP graft	7.0	26
BAS	M	64	Acute	ABF graft, ischemic heart disease, hypertension	-	Right SFA	3.0	6.0
TOU	M	68	Acute	Ischemic heart disease, hypertension	+	Right PA	1.0	6.0
Mean $\pm$ SEM		64 $\pm$ 4.1					7.8 $\pm$ 3.5	12 $\pm$ 4.0

ABF: aortobifemoral; AF: aortofemoral; CABG: coronary artery bypass grafting; CAD: coronary artery disease; CIA: common iliac artery; COPD: chronic obstructive pulmonary disease; DM: diabetes mellitus; EIA: external iliac artery; FF: femorofibular; FP: femoropopliteal; FT: femorotibial; IA: iliac artery; IF: iliofemoral; PA: popliteal artery; SFA: superficial femoral artery; TF: tibiofibular; AMI: acute myocardial infarction.

Table 16:

Treatment and outcome in patients with peripheral arterial occlusion, treated with SakSTAR(E65Q,K74Q,D82A,S84A,E108A,K109A,K130T,K135R,K136A,V137K), SakSTAR(K35A,E65Q,K74Q,D82A,S84A,T90A,E99D,T101S,E108A,K109A,K130T,K135R,K136A,V137K) or SakSTAR(K35A,E65Q,K74R,D82A,S84A,T90A,E99D,T101S,E108A,K109A,K130T,K135R,K136A,V137K)

Compound Patient Id.	Recanalization by thrombolysis	Total dose of thrombolytic agent (mg)	Total duration of infusion (hrs)	Additional therapy	Complications and remarks
SakSTAR(E65Q,K74Q,D82A,S84A,E108A,K109A,K130T,K135R,K136A,V137K) (SY118)					
VCL	Complete	17	22	None	Puncture site hematoma
REN	Complete	24	22	None	Puncture site hematoma
HOL	Complete	14	9.0	None	None
PAR	Partial (normal patency with residual thrombi after first control)	10	6.0		Small subdural hematoma
FRA	Complete	24	22	New FP graft	None
MAC	Complete	8.0	6.0	None	Brain stem hemorrhage; death
Mean $\pm$ SEM		16 $\pm$ 2.8	15 $\pm$ 3.4		
SakSTAR(K35A,E65Q,K74Q,D82A,S84A,T90A,E99D,T101S,E108A,K109A,K130T,K135R,K136A,V137K) (SY141)					
VERH	Complete	15	18	Femorofemoral cross over	
DUB	Complete	6.0	4.0		Puncture site hematoma
VAP	Complete	14	22	RIA-stenting, bilateral IA stenting	None
WYN	Complete	23	29	FP graft revision	None
HOR	Complete	10	8.0	None	Retroperitoneal hematoma, died due to septicemia
AND	Complete	13	17	None	
Mean $\pm$ SEM		14 $\pm$ 2.3	16 $\pm$ 3.7		
SakSTAR(K35A,E65Q,K74R,D82A,S84A,T90A,E99D,T101S,E108A,K109A,K130T,K135R,K136A,V137K) (SY145)					
LIN	Complete	14	24	None	Retroperitoneal hematoma. Hemorrhages
DEL	Complete	7.0	5.0	None	None
LAM	Complete	23	30	None	Puncture site hematoma
BAS	Complete			None	
TOU	Complete				
Mean $\pm$ SEM		15 $\pm$ 4.6	20 $\pm$ 7.5		

PTA, percutaneous transluminal angioplasty; IF, iliofemoral; FT, femoroiliac; FP, femoropopliteal.

Table 17: Neutralizing antibody activity before and after administration of SakSTAR(E65Q,K74Q,D82A,S84A,E108A,K109A,K130T,K135R,K136A,V137K), SakSTAR(K35A,E65Q,K74Q,D82A,S84A,T90A,E99D,T101S,E108A,K109A,K130T,K135R,K136A,V137K) or SakSTAR(K35A,E65Q,K74R,D82A,S84A,T90A,E99D,T101S,E108A,K109A,K130T,K135R,K136A,V137K) in patients with peripheral arterial occlusion

Neutralizing antibody activity (µg/ml)

Compound Patient Id.	Before	3 weeks	4 weeks
<b>SakSTAR(E65Q,K74Q,D82A,S84A,E108A,K109A,K130T,K135R,K136A,V137K) (SY118)</b>			
VCL	0.2	46	50
REN	0.1	1.6	0.9
HOL	0.2	22	18
PAR	0.1	19	15
FRA	1.2	15	39
MAC	0.0	-	-
Median	0.15	19	18
<b>SakSTAR(K35A,E65Q,K74Q,D82A,S84A,T90A,E99D,T101S,E108A,K109A,K130T,K135R,K136A,V137K) (SY141)</b>			
VERH	0.2	0.3	0.2
DUB	0.2	4.3	2.0
VAP	0.0	0.0	0.0
WYN	0.2	10	5.6
HOR	0.2	0.1	0.0
AND	0.8	-	100
Median	0.2	0.3	1.1
<b>SakSTAR(K35A,E65Q,K74R,D82A,S84A,T90A,E99D,T101S,E108A,K109A,K130T,K135R,K136A,V137K) (SY145)</b>			
LIN	0.0	5.5	3.8
DEL	0.2	1.6	1.9
LAM	0.2	164	190
BAS	0.1	-	-
TOU	-	-	-
Median	0.15	5.5	3.8

**Table 18: Immunogenicity of SakSTAR variants in patients with peripheral arterial occlusion**

	n	Neutralizing activity ( $\mu$ g/ml)	>5 $\mu$ g/ml	Specific IgG ( $\mu$ g/ml)	Code
SakSTAR	69	12 (4 - 100)	56	380 (81 - 1850)	.
SakSTAR(K74Q,E80A,D82A,K130T,K135R)	6	9.0 (0.1 - 23)	3	420 (31 - 730)	SY15
SakSTAR(E65D,K74R,E80A,D82A,K130T,K135R)	18	1.5 (0.2 - 7.0)	5	30 (24 - 100)	SY19
SakSTAR(E65Q,K74Q,D82A,S84A,E108A,K109A,K130T,K135R,K136A, $\nabla$ 137K)	6	27 (17 - 49)	5	2000 (1300 - 3600)	SY118
SakSTAR(K35A,E65Q,K74Q,D82A,S84A,T90A,E99D,T101S,E108A,K109A,K130T,K135R,K136A, $\nabla$ 137K)	6	0.7 (0.1 - 4.3)	2	7.7 (5.1 - 510)	SY141
SakSTAR(K35A,E65Q,K74R,D82A,S84A,T90A,E99D,T101S,E108A,K109A,K130T,K135R,K136A, $\nabla$ 137K)	3	4.7	1	.	SY145

80

Data represent median and 15-85 percentile range.

**Table 19: Cysteine substitution variants of SakSTAR**

Variant	Spec. Act. (kU/mg)	Dimerization level (%)	PEG derivatization	Clot lysis <i>in vitro</i> (C <sub>30</sub> in µg/ml)	t1/2(α) (min)	Clp (ml/min)	Antibody Absorption (Pool 40, %)
SakSTAR	130	0	none	0.33	2.0	2.2	95
SakSTAR (K102C)	143	0	none	0.29	nd	nd	95
SakSTAR (K102C-PEG)	108	0	1	0.60	3.0	0.32	
SakSTAR (K109C) monomeric	100	0	none	0.52	nd	nd	
SakSTAR (K109C) dimeric	1,650	>60	none	0.17	3.6	0.52	90
	2,235	>95	none	0.12	nd	nd	

**Table 20:** Cysteine-substitution variants of SakSTAR with reduced immunogenicity, substituted with maleimide-polyethylene glycol

Code	Fibrinolytic potency				Antibody absorption P40 (%)
	Specific activity (kU/mg) (C <sub>50</sub> : µg/ml)	Human plasma (C <sub>50</sub> : µg/ml)	Hamsters bolus (C <sub>50</sub> : µg/kg)	Cip (ml/min)	
		130	120		
SakSTAR					
	130	0.23	120	2.2	95
	140	0.24		3.7	57
SY19	51	0.37	42	0.45	58
SY19(S3C-SP5)*	20	0.65		0.28	50
SY19(S3C-MP5)*	43	0.42	20	0.15	57
SY19(S2C-SP5,S3C-SP5)	60	0.70	18	0.065	57
SY19(S3C-P20)	17	0.56	20	0.19	51
SY19(S3C-P10)					
	3,700	0.19		0.95	24
SY141	1,200	0.24	12		18
SY141(S3C-SP5)					18
SY141(S2C-SP5,S3C-SP5)	1,400	0.28			
	65	0.33	6	0.08	32
SY160(S3C-P20)	71	0.36	15	0.56	35
SY161(S3C-MP5)	66	0.40	9	0.15	38
SY161(S3C-P10)	155	0.32	8	0.04	44
SY161(S3C-P20)					

\*SP5: OPSS-PEG 5 kDa, MP5: MAL-PEG 5 kDa; P10: MAL-PEG 10 kDa; P20: MAL-PEG 20 kDa.

## REFERENCES

1. Lack CH: Staphylokinase: an activator of plasma protease. *Nature* 161: 559, 1948.
- 5 2. Lewis JH, Ferguson JH: A proteolytic enzyme system of the blood. III. Activation of dog serum profibrinolysin by staphylokinase. *Am J Physiol* 166: 594, 1951.
3. US5336495 (issued 94.08.09).
- 10 4. Vanderschueren S, Barrios L, Kerdsinchai P, Van den Heuvel P, Hermans L, Vrolix M, De Man F, Benit E, Muyldermans L, Collen D, Van de Werf F: A randomized trial of recombinant staphylokinase versus alteplase for coronary artery patency in acute myocardial infarction. 15 *Circulation* 92: 2044-2049, 1995.
5. Sako T, Sawaki S, Sakurai T, Ito S, Yoshizawa Y, Kondo I: Cloning and expression of the staphylokinase gene of *Staphylococcus aureus* in *Escherichia coli*. *Molec Gen Genet* 190: 271-277, 1983.
- 20 6. Behnke D, Gerlach D: Cloning and expression in *Escherichia coli*, *Bacillus subtilis*, and *Streptococcus sanguis* of a gene for staphylokinase - a bacterial plasminogen activator. *Molec Gen Genet* 210: 528-534, 1987.
- 25 7. Collen D, Silence K, Demarsin E, De Mol M, Lijnen HR: Isolation and characterization of natural and recombinant staphylokinase. *Fibrinolysis* 6: 203-213, 1992.
8. Sako T, Tsuchida N: Nucleotide sequence of 30 the staphylokinase gene from *Staphylococcus aureus*. *Nucleic Acids Res* 11: 7679-7693, 1983.
9. Collen D, Zhao ZA, Holvoet P, Marynen P: Primary structure and gene structure of staphylokinase. *Fibrinolysis* 6: 226-231, 1992.
- 35 10. Sakai M, Watanuki M, Matsuo O: Mechanism of fibrin-specific fibrinolysis by staphylokinase: participation of  $\alpha_2$ -plasmin inhibitor. *Biochem Biophys Res Comm* 162: 830-837, 1989.

007260-0640960



11. Matsuo O, Okada K, Fukao H, Tomioka Y, Ueshima S, Watanuki M, Sakai M: Thrombolytic properties of staphylokinase. *Blood* 76: 925-929, 1990.
12. Lijnen HR, Van Hoef B, De Cock F, Okada K, Ueshima S, Matsuo O, Collen D: On the mechanism of fibrin-specific plasminogen activation by staphylokinase. *J Biol Chem* 266: 11826-11832, 1991.
13. Lijnen HR, Van Hoef B, Matsuo O, Collen D: On the molecular interactions between plasminogen-staphylokinase,  $\alpha_2$ -antiplasmin and fibrin. *Biochim Biophys Acta* 1118: 144-148, 1992.
14. Silence K, Collen D, Lijnen HR: Interaction between staphylokinase, plasmin(ogen) and  $\alpha_2$ -antiplasmin. Recycling of staphylokinase after neutralization of the plasmin-staphylokinase complex by  $\alpha_2$ -antiplasmin. *J Biol Chem* 268: 9811-9816, 1993.
15. Silence K, Collen D, Lijnen HR: Regulation by  $\alpha_2$ -antiplasmin and fibrin of the activation of plasminogen with recombinant staphylokinase in plasma. *Blood* 82: 1175-1183, 1993.
16. Sakharov DV, Lijnen HR, Rijken DC. Interactions between staphylokinase, plasmin(ogen), and fibrin. *J Biol Chem* 271: 27912-27918, 1996.
17. Schlott B, Ghys KH, Hartmann M, Rcker A, Collen D. Staphylokinase requires  $\text{NH}_2$ -terminal proteolysis for plasminogen activation. *J Biol Chem* (in press).
18. Collen D, De Cock F, Vanlinthout I, Declerck PJ, Lijnen HR, Stassen JM. Comparative thrombolytic and immunogenic properties of staphylokinase and streptokinase. *Fibrinolysis* 6: 232-242, 1992.
19. Collen D, De Cock F, Stassen JM. Comparative immunogenicity and thrombolytic properties toward arterial and venous thrombi of streptokinase and recombinant staphylokinase in baboons. *Circulation* 87: 996-1006, 1993.
20. White H: Thrombolytic treatment for recurrent myocardial infarction. *Br Med J* 302: 429-430, 1991.

001260" 06470960

21. Gase A, Hartmann M, Ghys KH, Rcker A, Collen D, Behnke D, Schlott B: Functional significance of NH<sub>2</sub>- and COOH-terminal regions of staphylokinase in plasminogen activation. *Thromb Haemost* 76: 755-760, 1996.
- 5 22. EP 95200023.0 (January 6, 1995) and US 08/499,092 (July 6, 1995).
23. Schlott B, Hartmann M, Ghys KH, Birch-Hirschfeld E, Pohl HD, Vanderschueren S, Van de Werf F, Michoel A, Collen D, Behnke D: High yield  
10 production and purification of recombinant staphylokinase for thrombolytic therapy. *Bio/technology* 12: 185-189, 1994.
24. Horton RM, Hunt HD, Ho SN, Pullen JK, Pease LR. Engineering hybrid genes without the use of  
15 restriction enzymes: gene splicing by overlap extension. *Gene* 77: 61-68, 1989.
25. BIAcore system manual, 5-2, Pharmacia Biosensor AB, Uppsala, Sweden.
26. Karlsson R, Michaelsson A, Mattsson L:  
20 Kinetic analysis of monoclonal antibody-antigen interactions with a new biosensor based analytical system. *J Immunol Methods* 145: 229-240, 1991.
27. Sambrook J, Fritsch EF, Maniatis T: Molecular cloning: a laboratory manual. 2nd Ed. Cold Spring  
25 Harbor, NY. Cold Spring Harbor Laboratory Press, 1989.
28. Tartof KD, Hobbs CA: Improved media for growing plasmid and cosmid clones. *Bethesda Res Lab Focus* 9: 12, 1987
29. Bradford MM: A rapid and sensitive method  
30 for the quantitation of microgram quantities of protein utilizing the principle of protein-dye binding. *Anal Biochem* 72: 248, 1976.
30. Deutsch DG, Mertz ET: Plasminogen: purification from human plasma by affinity  
35 chromatography. *Science* 170: 1095-1096, 1970.
31. Collen D, Moreau H, Stockx L, Vanderschueren S. Recombinant staphylokinase variant with

007260"05470960

altered immunoreactivity. II. Thrombolytic properties and antibody induction. Circulation 94: 207-216, 1996.

32. Vanderschueren S, Stockx L, Wilms G, Lacroix H, Verhaeghe R, Vermylen J, Collen D:

- 5 Thrombolytic therapy of peripheral arterial occlusion with recombinant staphylokinase. Circulation 92: 2050-2057, 1995.

33. Gibaldi M, Perrier D. Pharmacokinetics, Marcel Dekker, New York, N.Y., 1983, 45-111.

- 10 34. Inada Y, Furukawa M, Sasaki H, Koderu Y, Hiroto M, Nishimura H, Matsushima A. Biomedical and biotechnological applications of PEG- and PM-modified proteins, TIBTECH 13: 86-91, 1995.

35. Collen D, Bernaerts R, Declerck P, De Cock  
15 F, Demarsin E, Jenn S, Laroche Y, Lijnen HR, Silence K, Verstrecken M. Recombinant staphylokinase variants with altered immunoreactivity. I. Construction and characterization. Circulation 94: 197-206, 1996.

36. Rabijns A, De Bondt HL, De Ranter C.  
20 Three-dimensional structure of staphylokinase, a plasminogen activator with therapeutic potential. Nature Struct Biol 4: 357-360, 1997.

007250 05470960

PCT/EP99/00748

Enclosure to letter dated 1 May 2000

87  
1

- 1. 05. 2000

(51)

## CLAIMS

1. Staphylokinase derivatives having essentially the amino acid sequence as depicted in figure 1 in which one or more amino acids have been replaced by another amino acid thus reducing the reactivity with a panel of murine monoclonal antibodies provided that the other amino acid is not alanine.

2. Staphylokinase derivatives as claimed in claim 1 having essentially the amino acid sequence as depicted in figure 1 in which one or more amino acids have been replaced by another amino acid thus reducing the absorption of SakSTAR-specific antibodies from plasma of patients treated with staphylokinase provided that the other amino acid is not alanine.

3. Staphylokinase derivatives as claimed in claim 1 having essentially the amino acid sequence as depicted in figure 1 in which one or more amino acids have been replaced by other amino acids, without reducing the specific activity by more than 50 percent provided that the other amino acid is not alanine.

4. Staphylokinase derivatives SakSTAR(K35X, G36X, E65X, K74X, E80X, D82X, K102X, E108X, K109X, K121X, K130X, K135X, K136X, +137X) having the amino acid sequence as depicted in figure 1 in which one or more of the amino acids Lys in position 35, Gly in position 36, Glu in position 65, Lys in position 74, Glu in position 80, Asp in position 82, Lys in position 102, Glu in position 108, Lys in position 109, Lys in position 121, Lys in position 130, Lys in position 135 and/or Lys in position 136 have been replaced with other amino acids provided that the other amino acid is not alanine and/or in which one amino acid has been added at the COOH-terminus, thus altering the immunogenicity after administration in patients, without markedly reducing the specific activity.

5. Staphylokinase derivatives listed in Tables 1, 3, 4, 5, 6, 7, 8, 13, 19 and 20, having the amino acid sequence as depicted in figure 1 in which the indicated

PCT/EP99/00748

88  
2

Enclosure to letter dated 1 May 2000

amino acids have been replaced by other amino acids thus reducing the absorption of SakSTAR-specific antibodies from plasma of patients treated with staphylokinase, without reducing the specific activity, provided that at least one amino acid is replaced with an amino acid other than alanine.

6. Staphylokinase derivative as claimed in claims 1-5 selected from the group consisting of SakSTAR(S34G,G36R,H43R), SakSTAR(S34G,G36R,H43R),
- 10 SakSTAR(G36R), SakSTAR(H43R), SakSTAR(G36R,K74R), SakSTAR(K35E), SakSTAR(K74Q), SakSTAR(K130T), SakSTAR(V132L), SakSTAR(V132T), SakSTAR(V132N), SakSTAR(V132R), SakSTAR(K130T,K135R), SakSTAR(G36R, K130T,K135R), SakSTAR(K74R,K130T,K135R), SakSTAR(K74Q, K130T,K135R), SakSTAR(G36R,K74R,K130T,K135R),
- 15 SakSTAR(G36R,K74Q,K130T,K135R), SakSTAR(G36R,H43R,K74R, K130T,K135R), SakSTAR(E65A,K74Q,K130T,K135R), SakSTAR(E65Q,K74Q,K130T,K135R), SakSTAR(K74Q,K86A,K130T, K135R), SakSTAR(E65Q,T71S,K74Q,K130T,K135R),
- 20 SakSTAR(K74Q,K130A,K135R), SakSTAR(E65Q,K74Q,K130A, K135R), SakSTAR(K74Q,K130E,K135R), SakSTAR(K74Q,K130E, V132R,K135R), SakSTAR(E65Q,K74Q,T90A,K130A,K135R), SakSTAR(E65Q,K74Q,N95A,K130A,K135R), SakSTAR(E65Q,K74Q, E118A,K130A,K135R), SakSTAR(E65Q,K74Q,N95A,E118A,K130A, K135R), SakSTAR(N95A,K130A,K135R), SakSTAR(E65Q,K74Q, K109A,K130,K135R), SakSTAR(E65Q,K74Q,E108A,K109A, K130T,K135R), SakSTAR(E65Q,K74Q,K121A,K130T,K135R),
- 25 SakSTAR(E65Q,K74Q,N95A,E118A,K130A,K135R,K136A,+137K), SakSTAR(E80A,D82A,K130T,K135R), SakSTAR(K74R,E80A,D82A, K130T,K135R), SakSTAR(K74Q,E80A,D82A,K130T,K135R), Sak- STAR(K35A,K74R,E80A,D82A,K130T,K135R), SakSTAR(E65D, K74R,E80A,D82A,K130T,K135R), SakSTAR(E65S,K74R,E80A, D82A,K130T,K135R), SakSTAR(S34G,G36R,K74R,K130T,K135R), SakSTAR(E65A,K74R,E80A,D82A,K130T,K135R), SakSTAR(E65N, K74R,E80A,D82A,K130T,K135R), SakSTAR(E65Q,K74R,E80A, D82A,K130T,K135R), SakSTAR(K57A,E58A,E61A,E80A,D82A, K130T,K135R), SakSTAR(E65D,K74Q,E80A,D82A,K130T,K135R), SakSTAR(E65Q,K74Q,E80A,D82A,K130T,K135R), SakSTAR(K35A,

PCT/EP99/00748

89

Enclosure to letter dated 1 May 2000

E65D, K74Q, E80A, D82A, K130T, K135R), SakSTAR (K74R, E80A, D82A, S103A, K130T, K135R), SakSTAR (E65D, K74R, E80A, D82A, K109A, K130T, K135R), SakSTAR (E65D, K74R, E80A, D82A, K130T, K135R, K136A), SakSTAR (E65Q, K74Q, D82A, S84A, K130T, K135R),  
 5 SakSTAR (K35A, K74Q, E80A, D82A, K130T, K135R), SakSTAR (K35A, E65D, K74R, E80A, D82A, K130T, K135R).

7. SakSTAR (E65D, K74R, E80A, D82A, K130T, K135R) having the code SY19.

8. SakSTAR (K35A, E65Q, K74R, E80A, D82A, T90A, E99D, T101S, E108A, K109A, K130T, K135R) having the code SY161.

9. Staphylokinase derivatives having essentially the amino acid sequence as depicted in figure 1 in which one or more amino acids have been replaced by another amino acid thus reducing the reactivity with a  
 15 panel of murine monoclonal antibodies and having in addition either one or both of the following:

- at least one amino acid substituted with Cys, resulting in dimerization and/or increased specific activity and/or reduced clearance and/or increased throm-  
 20 bolytic potency; and/or

- polyethylene glycol substitution, resulting in a significantly reduced plasma clearance while maintaining specific activity.

10. Staphylokinase derivatives as claimed in  
 25 claim 9 having essentially the amino acid sequence as depicted in figure 1 in which one or more amino acids have been replaced by another amino acid thus reducing the absorption of SakSTAR-specific antibodies from plasma of patients treated with staphylokinase.

30 11. Staphylokinase derivatives as claimed in claim 9 having essentially the amino acid sequence as depicted in figure 1 in which one or more amino acids have been replaced by other amino acids, without reducing the specific activity by more than 50 percent.

35 12. Staphylokinase derivatives as claimed in claim 9, named SakSTAR (K35X, G36X, E65X, K74X, E80X, D82X, K102X, E108X, K109X, K121X, K130X, K135X, K136X, +137X) and having the amino acid sequence as depicted in figure 1 in which

PCT/EP99/00748

90  
4

Enclosure to letter dated 1 May 2000

one or more of the amino acids Lys in position 35, Gly in position 36, Glu in position 65, Lys in position 74, Glu in position 80, Asp in position 82, Lys in position 102, Glu in position 108, Lys in position 109, Lys in position 121, Lys in position 130, Lys in position 135 and/or Lys in position 136 have been replaced with other amino acids and/or in which one amino acid has been added at the COOH-terminus, thus altering the immunogenicity after administration in patients, without markedly reducing the specific activity.

13. Staphylokinase derivatives as claimed in claim 9 and listed in Tables 1, 3, 4, 5, 6, 7, 8, 13, 19 and 20, having the amino acid sequence as depicted in figure 1 in which the indicated amino acids have been replaced by other amino acids thus reducing the absorption of SakSTAR-specific antibodies from plasma of patients treated with staphylokinase, without reducing the specific activity.

14. Staphylokinase derivative as claimed in claims 9-13 selected from the group consisting of SakSTAR(K74A,E75A,R77A), SakSTAR(K35A,E75A), SakSTAR(E75A), SakSTAR(E80A,D82A), SakSTAR(E80A), SakSTAR(D82A), SakSTAR(E75A,D82A), SakSTAR(S34G,G36R,H43R), SakSTAR(K35A), SakSTAR(D82A), SakSTAR(D82A,S84A), SakSTAR(T90A), SakSTAR(Y92A), SakSTAR(K130A), SakSTAR(V132A), SakSTAR(S34G,G36R,H43R), SakSTAR(G36R), SakSTAR(H43R), SakSTAR(G36R,K74R), SakSTAR(K35E), SakSTAR(K74Q), SakSTAR(K130T), SakSTAR(V132L), SakSTAR(V132T), SakSTAR(V132N), SakSTAR(V132R), SakSTAR(K130T,K135R), SakSTAR(G36R,K130T,K135R), SakSTAR(K74R,K130T,K135R), SakSTAR(K74Q,K130T,K135R), SakSTAR(G36R,K74R,K130T,K135R), SakSTAR(G36R,K74Q,K130T,K135R), SakSTAR(G36R,H43R,K74R,K130T,K135R), SakSTAR(E65A,K74Q,K130T,K135R), SakSTAR(E65Q,K74Q,K130T,K135R), SakSTAR(K74Q,K86A,K130T,K135R), SakSTAR(E65Q,T71S,K74Q,K130T,K135R), SakSTAR(K74Q,K130A,K135R), SakSTAR(E65Q,K74Q,K130A,K135R), SakSTAR(K74Q,K130E,K135R), SakSTAR(K74Q,K130E,

PCT/EP99/00748

91  
5

Enclosure to letter dated 1 May 2000

V132R, K135R), SakSTAR (E65Q, K74Q, T90A, K130A, K135R), Sak-  
 STAR (E65Q, K74Q, N95A, K130A, K135R), SakSTAR (E65Q, K74Q,  
 E118A, K130A, K135R), SakSTAR (E65Q, K74Q, N95A, E118A, K130A,  
 K135R), SakSTAR (N95A, K130A, K135R), SakSTAR (E65Q, K74Q,  
 5 K109A, K130, K135R), SakSTAR (E65Q, K74Q, E108A, K109A,  
 K130T, K135R), SakSTAR (E65Q, K74Q, K121A, K130T, K135R),  
 SakSTAR (E65Q, K74Q, N95A, E118A, K130A, K135R, K136A, +137K),  
 SakSTAR (E80A, D82A, K130T, K135R), SakSTAR (K74R, E80A, D82A,  
 K130T, K135R), SakSTAR (K74Q, E80A, D82A, K130T, K135R), Sak-  
 10 STAR (K35A, K74R, E80A, D82A, K130T, K135R), SakSTAR (E65D,  
 K74R, E80A, D82A, K130T, K135R), SakSTAR (E65S, K74R, E80A,  
 D82A, K130T, K135R), SakSTAR (S34G, G36R, K74R, K130T, K135R),  
 SakSTAR (E65A, K74R, E80A, D82A, K130T, K135R), SakSTAR (E65N,  
 K74R, E80A, D82A, K130T, K135R), SakSTAR (E65Q, K74R, E80A,  
 15 D82A, K130T, K135R), SakSTAR (K57A, E58A, E61A, E80A, D82A,  
 K130T, K135R), SakSTAR (E65D, K74Q, E80A, D82A, K130T, K135R),  
 SakSTAR (E65Q, K74Q, E80A, D82A, K130T, K135R), SakSTAR (K35A,  
 E65D, K74Q, E80A, D82A, K130T, K135R), SakSTAR (K74R, E80A, D82A,  
 S103A, K130T, K135R), SakSTAR (E65D, K74R, E80A, D82A, K109A,  
 20 K130T, K135R), SakSTAR (E65D, K74R, E80A, D82A, K130T,  
 K135R, K136A), SakSTAR (E65Q, K74Q, D82A, S84A, K130T, K135R),  
 SakSTAR (K35A, K74Q, E80A, D82A, K130T, K135R), SakSTAR (K35A,  
 E65D, K74R, E80A, D82A, K130T, K135R).

15. Staphylokinase derivatives as claimed in  
 25 claim 9 wherein the Cys is chemically modified with  
 polyethylene glycol with molecular weights up to 20 kDa.

16. Staphylokinase derivatives as claimed in  
 claim 15 wherein selected amino acids in the NH<sub>2</sub>-terminal  
 region of 10 amino acids, are substituted with Cys, which  
 30 is chemically modified with polyethylene glycol with  
 molecular weights up to 20 kDa, which derivatives are  
 characterized by a significantly reduced plasma clearance  
 and maintained thrombolytic potency upon single intrave-  
 nous bolus administration at a reduced dose.

35 17. Staphylokinase derivative as claimed in  
 claim 16, wherein the serine in position 2 or 3 is  
 substituted with a cystein and the cystein is chemically



PCT/EP99/00748

92.

Enclosure to letter dated 1 May 2000

modified with polyethylene glycol having a molecular weight of 5, 10 or 20 kDa.

18. Staphylokinase derivative as claimed in claim 17, which derivative is SakSTAR(S3C-MP5,K35A,E65Q, 5 K74R,E80A,D82A,T90A,E99D,T101S,E108A,K109A,K130T,K135R) (SY161(S3C-MP5)).

19. Staphylokinase derivative as claimed in claim 17, which derivative is SakSTAR(S3C-P10,K35A, E65Q,K74R,E80A,D82A,T90A,E99D,T101S,E108A,K109A,K130T, 10 K135R) (SY161(S3C-P10)).

20. Staphylokinase derivative as claimed in claim 17, which derivative is SakSTAR(S3C-P20,K35A, E65Q,K74R,E80A,D82A,T90A,E99D,T101S,E108A,K109A,K130T, K135R) (SY161(S3C-P20)).

15 21. Staphylokinase derivative as claimed in claim 17, which derivative is SakSTAR(S3C-MP5,E65D,K74R, E80A,D82A,K130T,K135R) (SY19(S3C-MP5)).

22. Staphylokinase derivative as claimed in claim 17, which derivative is SakSTAR(S3C-SP5,E65D,K74R, 20 E80A,D82A,K130T,K135R) (SY19(S3C-SP5)).

23. Staphylokinase derivative as claimed in claim 17, which derivative is SakSTAR(S2C-SP5,S3C-SP5, E65D,K74R,E80A,D82A,K130T,K135R) (SY19(S2C-SP5,S3C-SP5)).

24. Staphylokinase derivative as claimed in 25 claim 17, which derivative is SakSTAR(S3C-P20,E65D,K74R, E80A,D82A,K130T,K135R) (SY19(S3C-P20)).

25. Staphylokinase derivative as claimed in claim 17, which derivative is SakSTAR(S3C-P20,E65D,K74R, E80A,D82A,K130T,K135R) (SY19(S3C-P10)).

30 26. Dimer of two staphylokinase derivatives as claimed in claim 9.

27. Method for producing the staphylokinase derivatives as claimed in claims 1 to 8, comprising the steps of:

35 a. preparing a DNA fragment comprising at least the part of the coding sequence of staphylokinase that provides for its biological activity;

PCT/EP99/00748

7<sup>93</sup>

Enclosure to letter dated 1 May 2000

b. performing in vitro site-directed mutagenesis on the DNA fragment to replace one or more codons for wild-type amino acids by a codon for another amino acid;

c. cloning the mutated DNA fragment in a suitable vector;

d. transforming or transfecting a suitable host cell with the vector; and

e. culturing the host cell under conditions suitable for expressing the DNA fragment.

10 28. Method as claimed in claim 27, wherein the DNA fragment is a 453 bp EcoRI-HindIII fragment of the plasmid pMEX602sakB, the in vitro site-directed mutagenesis is performed and the mutated DNA fragment is expressed in E. coli.

15 29. Pharmaceutical composition comprising at least one of the staphylokinase derivatives as claimed in claims 1 to 25 together with a suitable excipient.

30. Pharmaceutical composition as claimed in claim 29 for treating arterial thrombosis.

IDENTIFICATION, PRODUCTION AND USE OF STAPHYLOKINASE  
DERIVATIVES WITH REDUCED IMMUNOGENICITY AND/OR REDUCED  
CLEARANCE

ABSTRACT OF THE DISCLOSURE

Methods for the identification, production and use  
of staphylokinase derivatives characterized by a reduced  
immunogenicity after administration in patients and that can  
be administered by single intravenous bolus injection. The  
5 derivatives of the invention are obtained by preparing a DNA  
fragment comprising at least the part of the coding sequence  
of staphylokinase that provides for its biological activity;  
performing *in vitro* site-directed mutagenesis on the DNA  
10 fragment to replace one or more codons for wild-type amino  
acids by a codon for another amino acid; cloning the mutated  
DNA fragment in a suitable vector; transforming or  
transfecting a suitable host cell with the vector; culturing  
the host cell under conditions suitable for expressing the  
15 DNA fragment; purifying the expressed staphylokinase  
derivative to homogeneity and chemically modifying  
substituted Cys residues with thiol-directed polyethylene  
glycol; preferably the DNA fragment is a 453 bp EcoRI-  
HindIII fragment of the plasmid pMEX602sakB, (pMEX.SakSTAR),  
the *in vitro* site-directed mutagenesis is performed by  
20 spliced overlap extension polymerase chain reaction and the  
mutated DNA fragment is expressed in *E. coli* strain TG1 or  
WK6. The invention also relates to pharmaceutical  
compositions comprising at least one of the staphylokinase  
derivatives according to the invention together with a  
25 suitable excipient, for treatment of arterial thrombosis.


1 14  
 Ser Ser Ser Phe Asp Lys Gly Lys Tyr Lys Lys Gly Asp Asp  
 15 28  
 Ala Ser Tyr Phe Glu Pro Thr Gly Pro Tyr Leu Met Val Asn  
 29 42  
 Val Thr Gly Val Asp Ser Lys Gly Asn Glu Leu Leu Ser Pro  
 43 56  
 His Tyr Val Glu Phe Pro Ile Lys Pro Gly Thr Thr Leu Thr  
 57 70  
 Lys Glu Lys Ile Glu Tyr Tyr Val Glu Trp Ala Leu Asp Ala


1/8

FIG. 1-I

71	Thr	Ala	Tyr	Lys	Glu	Phe	Arg	Val	Val	Glu	Leu	Asp	Pro	Ser	84
85	Ala	Lys	Ile	Glu	Val	Thr	Tyr	Tyr	Asp	Lys	Asn	Lys	Lys	Lys	98
99	Glu	Glu	Thr	Lys	Ser	Phe	Pro	Ile	Thr	Glu	Lys	Gly	Phe	Val	112
113	Val	Pro	Asp	Leu	Ser	Glu	His	Ile	Lys	Asn	Pro	Gly	Phe	Asn	126
127	Leu	Ile	Thr	Lys	Val	Val	Ile	Glu	Lys	Lys					136

FIG. 1-II










FIG. 3-II




6/8

FIG. 4-I

7/8

C



D





FIG. 4-II



# Declaration and Power of Attorney For Patent Application

## English Language Declaration

As a below named inventor, I hereby declare that:

My residence, post office address and citizenship are as stated below next to my name,

I believe I am the original, first and sole inventor (if only one name is listed below) or an original, first and joint inventor (if plural names are listed below) of the subject matter which is claimed and for which a patent is sought on the invention entitled Identification, production, and use of staphylokinase derivatives with reduced immunogenicity and/or reduced clearance, the specification of which

(check one)

☐ is attached hereto.

☒ was filed on February 4, 1999 as PCT international application ~~xxx~~

~~Application No.~~ PCT/EP99/00748 and 09/601,490 on 03 August 2000

and was amended on August 3, 2000

(if applicable)

I hereby state that I have reviewed and understand the contents of the above identified specification, including the claims, as amended by any amendment referred to above.

I acknowledge the duty to disclose information which is material to patentability of this application in accordance with Title 37, Code of Federal Regulations, §1.56.

I hereby claim foreign priority benefits under Title 35, United States Code, §119 of any foreign application(s) for patent or inventor's certificate listed below and have also identified below any foreign application for patent or inventor's certificate having a filing date before that of the application on which priority is claimed:

Prior Foreign Application(s)			Priority Claimed	
<u>98200323.8</u>	<u>Europe</u>	<u>February 4, 1998</u>	<input checked="" type="checkbox"/>	<input type="checkbox"/>
(Number)	(Country)	(Day/Month/Year Filed)	Yes	No
<u>98200365.9</u>	<u>Europe</u>	<u>February 6, 1998</u>	<input checked="" type="checkbox"/>	<input type="checkbox"/>
(Number)	(Country)	(Day/Month/Year Filed)	Yes	No
<u>                    </u>	<u>                    </u>	<u>                    </u>	<input type="checkbox"/>	<input type="checkbox"/>
(Number)	(Country)	(Day/Month/Year Filed)	Yes	No

I hereby claim the benefit under Title 35, United States Code, §120 of any United States application(s) listed below and, insofar as the subject matter of each of the claims of this application is not disclosed in the prior United States application in the manner provided by the first paragraph of Title 35, United States Code, §112, I acknowledge the duty to disclose material information as defined in Title 37, Code of Federal Regulations, §1.56(a) which occurred between the filing date of the prior application and the national or PCT international filing date of this application:

(Application Serial No.)	(Filing Date)	(Status) (patented, pending, abandoned)
--------------------------	---------------	--

(Application Serial No.)	(Filing Date)	(Status) (patented, pending, abandoned)
--------------------------	---------------	--

I hereby declare that all statements made herein of my own knowledge are true and that all statements made on information and belief are believed to be true; and further that these statements were made with the knowledge that willful false statements and the like so made are punishable by fine or imprisonment, or both, under Section 1001 of Title 18 of the United States Code and that such willful false statements may jeopardize the validity of the application or any patent issued thereon.

POWER OF ATTORNEY: As a named inventor, I hereby appoint the following attorney(s) and/or agent(s) to prosecute this application and transact all business in the Patent and Trademark Office connected therewith. (list name and registration number)

William H. Logsdon	22,132	Paul M. Reznick	33,059	Jesse A. Hirshman	40,016
Russell D. Orkin	25,363	John W. McIlvaine	34,219	James G. Porcelli	33,757
David C. Hanson	23,024	Michael I. Shamos	30,424	Kent E. Baldauf, Jr.	36,082
Richard L. Byrne	28,498	Blynn L. Shideler	35,034	Christian Schuster	43,908
Frederick B. Ziesenheim	19,438	Julie W. Meder	36,216	Dean E. Geibel	42,570
Kent E. Baldauf	25,826	Lester N. Fortney	38,141	Thomas J. Clinton	40,561
Barbara E. Johnson	31,198	Randall A. Notzen	36,882	Nathan J. Prepelka	43,016

Send Correspondence to:

Barbara E. Johnson, 700 Koppers Building, 436 Seventh Avenue, Pittsburgh PA 15219-1818

Direct Telephone calls to: (name and telephone number) Barbara E. Johnson (412) 471-8815

Full name of sole or first inventor

Collen, Désiré José

Inventor's signature

Date

Residence

Winkelse-Herent, Belgium BEX

Citizenship

Belgium

Post Office Address

Schoonzichtlaan 20, B-3020 Winkelse-Herent, Belgium

Full name of second joint inventor, if any

Second inventor's signature

Date

Residence

Citizenship

Post Office Address

(Supply similar information and signature for third and subsequent joint inventors.)